



## SCREENING OF SECONDARY METABOLITES AND ANTIBACTERIAL ACTIVITY OF *ACACIA CONCINNA* LEAVES

Xavier Vergeese raja\*, Rajeshwari Sivaraj

Department of Biotechnology, School of Life science, Karpagam University, Coimbatore, Tamilnadu, India

Article Received on: 18/08/12 Revised on: 12/09/12 Approved for publication: 05/10/12

\*Email:xavierbt\_06@yahoo.co.in

### ABSTRACT

The preliminary phytochemical study and *in vitro* antibacterial activity of the leaf extracts of *Acacia concinna* having ethno medicinal uses were investigated. The preliminary phytochemical analysis of the extracts revealed the presence of phenol, tannin, fat and fixed oil, flavanoids, saponin and quinone. The extracts were subjected for screening of *in vitro* antibacterial activity against various pathogens at the concentrations of 500 and 250 µg/ml by well diffusion assay method. The results of antibacterial activity revealed that all the extract showed good inhibitory activity against all the tested pathogens and the chloroform extract showed comparative by better activity than the other extracts against *S.pyogens* and *s. aureus*. The activities of the extract were compared with standard antibiotics. These results indicate that *A. concinna* leaves possesses potential broad spectrum antibacterial activity.

**KEYWORDS:** *Acacia concinna*, phytochemical screening, bioactive compounds, herbal remedies.

### INTRODUCTION

Knowledge of the chemical constituent of plants is desirable for the discovery of therapeutic agents and in discovering the actual value of folklore remedies. Traditionally, screening methods have been used to study the pharmacological effects of phytochemical compounds. The medicinal value of these plants lies in some chemical substances that produce a definite physiological action on the human body<sup>1</sup>.

The most important of these bioactive constituents of plants are alkaloids, tannins, flavonoids and phenolic compounds. Biologically active compounds. from natural sources have always been of great interest to scientists working on infectious diseases. World Health Organization advocated that both developed and developing countries should interact with the traditional medicine with a view of safe and effective remedies of ailments.

*Acacia concinna* is a tree native of Asia. Its parts used as bark, leaves and pods. It is a common, prickly, scandent shrub, occurring in tropical jungles throughout India, especially in the Deccan. An infusion of the leaves is used in malarial fever<sup>2</sup>. A decoction of the pods relieves biliousness and acts as a purgative. The pods are reported to be used in north Bengal for poisoning fish<sup>3</sup>. The present study investigates the presence of secondary metabolites and antimicrobial activities against some human pathogens.

### MATERIALS AND METHODS

#### Sample Collection and Preparation

*Acacia concinna* leaves were collected from Coimbatore district, Tamilnadu, India. The plant leaves selected for the study were washed, air dried and powdered. 250 g of *Acacia concinna* leaves yields 200 g powder.

#### Preparation of The Extract

About 50 g of dried powder samples were weighed and extraction process was carried out by using 200 ml of solvents (Ethanol, Methanol and Chloroform) in soxhlet apparatus for 18 hours. The extract was concentrated by evaporation at 100° C for 8 hours and then air dried. The concentrated extract were made in to a fine powder form and stored at room temperature prior to phytochemical screening.

#### Phytochemical and Biochemical Analysis

The phytochemical tests were carried out using different solvents extracts using standard procedures to identify the constituents<sup>4</sup>. To assess the activity of selected medicinal plants, preliminary phytochemical analysis studies were carried out.

#### Bacterial Cultures

Eleven different bacterial cultures were tested for the activity of the extract.

#### Preparation of Bacterial culture

The stock culture of the four bacteria used was sub-cultured on Nutrient Agar at 37° C for 24 hours. The culture was emulsified in sterile saline.

#### Sensitivity Test

The agar diffusion method was used. Wells (7mm) were cut by using gel puncture and the previously prepared cultured saline was swabbed on the culture plates containing Nutrient Agar plates. The same quantity of sterile water and solvents used acts as the negative control. A volume of 50µl plant sample with different extracts were tested in a concentration of 100 mg/ml and incubation was performed at 37° C for 24 hours. The assessment of antimicrobial activity was based on the measurement of the diameter of the inhibition zone formed around the well. Standardized streptomycin was used as the antibiotic control (Positive control). Triplicate plates were prepared for each extract and controls.

**Table 1: Phytochemical screening of *Acacia concinna* leaves**

S.No	Reagent	Nature of colour change	Inference	Phytochemical changes
1.	Substance + alcohol + FeCl <sub>3</sub>	Greenish yellow	Present	Presence of phenol
2.	0.5g substance + 20mL H <sub>2</sub> O is boiled. Then 0.1% FeCl <sub>3</sub>	Brownish green	Present	Presence of tannin
3.	Substance + Sudan III	Shining orange colour	Present	Presence of fat and fixed oil
4.	Substance + 10% NaOH	Green brown	Present	Presence of flavonoids
5.	Substance shaken in water	Frothing present	Present	Presence of saponin
6.	Substance + chloroform+ drop of acetic acid, heated + conc. H <sub>2</sub> SO <sub>4</sub>	Orange	Absent	Absence of steroids
7.	Substance + conc. HCl	Green	Present	Presence of quinone
8.	Substance + Iodine followed by H <sub>2</sub> SO <sub>4</sub>	Brown	Absent	Absence of cellulose
9.	Substance + 2 mL chloroform + conc. H <sub>2</sub> SO <sub>4</sub>	Light orange	Absent	Absence of terpenoids
10.	Substance + 2 mL glacial acetic acid + 1 drop of FeCl <sub>3</sub> + 1 mL conc. H <sub>2</sub> SO <sub>4</sub>	Brown	Absent	Absence of glycosides

**Table 2: Antibacterial activity of leaf extract of *Acacia concinna* (Leaves)**

organism	Ethanol (cm)				Methanol (cm)				Chloroform (cm)			
	A(20µl)	B(40µl)	C(60µl)	D(100µl)	A	B	C	D	A	B	C	D
<i>P.aerogenosa</i>	0.7	1	1.8	2	0.2	0.4	0.8	1	0.3	0.7	1.2	1.5
<i>P.nirabilis</i>	0.2	0.5	0.6	1	0.2	0.6	0.9	1.2	0.1	0.6	1.1	1.5
<i>Str.pyogens</i>	0.3	0.5	0.8	1.4	0.1	0.5	1.2	2	1.3	1.6	1.7	2.5
<i>E.coli</i>	0.4	0.8	1.2	1.9	0.1	0.3	0.5	0.8	0.5	1.2	1.6	2
<i>Yersinia</i>	0.3	0.6	1.2	1.8	0.4	0.6	1.2	1.4	0.7	1.1	1.7	2.1
<i>E.aerogenes</i>	0.5	0.8	1.5	1.8	0.2	0.4	0.9	1.8	0.4	1.1	1.5	2
<i>S.typhi</i>	0.2	0.3	0.5	0.7	0.1	0.3	0.6	0.8	0.1	0.3	0.5	0.9
<i>E.aerogenes</i>	0.1	0.5	1	1.7	0.5	0.9	1.2	1.5	0.9	1.3	1.4	1.8
<i>S.aureus</i>	0.1	0.4	0.8	1.2	1	1.8	2	2.3	0.5	1.4	2.1	2.4
<i>K.pneumonia</i>	0.1	0.4	0.7	1	0.3	0.5	1	1.2	0.3	0.5	0.8	1.5
<i>B.subtilis</i>	0.2	0.6	1.2	1.7	0.4	0.6	0.9	1.3	0.2	0.8	1.1	1.9

## RESULTS

The phytochemical screening of the test plant was done for their active components present for their medicinal values. Many of the phytochemical analysis showed positive results which render the presence of their active compounds like phenol, tannin, fat and fixed oil, flavanoids, saponin and quinone.

The extracts of *A.concinna* obtained using different solvents were effective against the bacterial strains (Table 02). Among different extracts Chloroform exhibited maximum antimicrobial activity. The increased zone on inhibition was measured in *S.pyogens* (25 mm) compared to other microorganisms, which was followed by *S. aureus* (24 mm).

## DISCUSSION

The ethanol extracts of *Balanites aegyptica* leaves, stem bark, root bark and fruit were tested for antimicrobial activities *Bacillus subtilis*, *Staphylococcus aureus*, *E.coli*, *Pseudomonas aeruginosa* and *Candida albicans* and the maximum inhibition zone was recorded in the leaf extract<sup>5</sup>.

The essential from *Aframomum danielli*, *Zingiber officinale* and *Curcuma longa* have antimicrobial activity against both

gram positive and gram negative bacteria, yeast and filamentous fungi<sup>6</sup>.

The water and alcoholic extracts of leaves of *Melia dubia* when tested in gram positive and gram negative strains using agar gel diffusion method exhibited moderately good antimicrobial activity<sup>7</sup>

## REFERENCES

- Parekh J, Darshana J and Sumitra C. Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. *Journal of Ecology*, 2005, 29: 203-210.
- Krishnaiah D, Sarbatly R, Bono A. Phytochemical antioxidants for health and medicine - A move towards nature. *Biotechnol. Mol. Biol. Rev* 2007; 1(4):097-104.
- Singh SP and Mishra N. An experimental Study of analgesic activity of *Cissus quadrangularis*. *Ind. J. Pharmacol.* 1984: 167.
- Harborne JB. "Phytochemical methods", Chapman and Hall, Ltd. London, 1999; P.60-66.
- Augusti KT and Cherian S. Insulin sparing action of leucopelargonidin derivative isolated from *Ficus bengalensis* Linn. *Indian J. Exp. Biol.* 2008, 33: 608-611.
- Udayakumar RH, Velmurugan D, Sivanesan and Krishna R. Phytochemical and antimicrobial studies of extracts of *Solanum xanthocarpum*. *Ancient Sci. Life*, 2003; 23(2).p.90.
- Naseer Banu A, Sivakumar, Subramanian MS. Effect of manures on biomass production and pharmaco-biochemical properties of some greens. *Ancient Sci. Life*, 2003, 23(2).p.123-130.

Source of support: Nil, Conflict of interest: None Declared