



## Review Article

### A NOVEL REVIEW ON *ERYTHRINA VARIEGATA*

Lahari K\*, Divya M, Vidyavathi N, Kishore L, Poojitha M

Department of Pharmacology, Sree Vidyanikethan College of Pharmacy, Sree Sainath Nagar, A Rangampet, Chittoor District, Andhra Pradesh, India

\*Corresponding Author Email: lallu.sahana@gmail.com

Article Received on: 01/04/15 Revised on: 23/04/15 Approved for publication: 28/04/15

**DOI: 10.7897/2230-8407.06451**

#### ABSTRACT

Medicinal plants are nature's gift to human society to make disease free healthy life. More than thousand medicinal plants are recognised in our country. The present review is therefore an effort to give a detail survey of the literature on its phytopharmacological properties of *Erythrina variegata* belonging to the family Fabaceae, which is a shrub with prickly stems; it is a wild growing forest plant in India. Majorly popular system of medicine like Ayurveda, Siddha, Unani and homeopathy. Various plant parts such as bark, root, fruits and leaves are used in treatment of fever, astringent, febrifuge, skin diseases etc.

**Keywords:** *Erythrina variegata*, Fabaceae, Haematological parameters, Green medicine.

#### INTRODUCTION

Medicinal plants are majorly an important therapeutic aid for alleviating the ailments of human kind. The current widespread and strong belief about the plant derived drugs is that "Green medicine" is safe and more dependable than the costly synthetic drugs due to their adverse side effects.

#### Description

The genus *Erythrina* comprises majorly comprises of about 110 species of trees. Here, the name "coral tree" is used as a collective term for these plants. *Erythrina variegata* is a medium sized deciduous small tree having prickly stems and branches, leaves having triangular leaflets and with large coral red flowers.

Majorly grows over Bangladesh and almost spread widely throughout India. It has been worked as a folk medicine in the tropical and sub-tropical regions; it is known to possess pharmacological activities. This most attractive type of plant is majorly grown for its variegated leaves and also for its showy red flowers. This is a fast growing plant, grows up to 50-60 feet tall and having green and yellow leaves about 6 inches long, during spring season this coral tree is decorated with showy red blossoms of 2.5 inches, which are arranged in dense, 6 inches long racemes. These blooms are followed by 12 inch long red/brown seed pods which contain poisonous seeds.

*Erythrina* generally is a Greek word "Eruthros" which means red and it shows red species<sup>1,2</sup>.

Telugu : Badisa, Baditha, Kalyana murungi  
Kingdom: Plantae  
(unranked): Angiosperms  
Class: Magnoliopsida – Dicotyledons  
Order: Fabales  
Family: Fabaceae  
Subfamily: Papilionoideae  
Genus: *Erythrina*

Species: *variegata*

#### Morphology

Plant height: 50 – 60 feet  
Plant type: Thorny tree  
Stem type: Soft, smooth, prickly stem  
Leaf type : Trifoliate (6 inches)  
Leaf color: Green  
Flower : Dark red (2.5 inches)  
Odour : Pungent  
Bark : Brownish grey

#### Uses

*Erythrina variegata* is used as a wind break and shade tree with in common plants.

The bark of the plant is astringent, febrifuge, antibilious and antihelmintic. This is also used in ophthalmic and various skin diseases, nervine sedative, febrifuge, anti-asthmatic and anti epileptic.

The leaves are used in fever, inflammation and joint pain. They have a potential use in the prevention of postmenopausal bone loss. It is also known to possess pharmacological activities including antimicrobial, anti bacterial effects.

The juice of the leaves is mixed with honey and ingested to kill tapeworm, round worm and threadworm. It is also useful in ophthalmic and skin diseases<sup>2,3</sup>.

#### Chemical constituents

The bark and root of *Erythrina variegata* Linn showed the presence of carbohydrates, glycosides, proteins, volatile oils and tannins. The plant is a rich source of alkaloids (2.5%). In this flavanoids are chemical phenyl benzopyrones, usually conjugated with sugars are

present in all vascular plants. Isoflavonoids are reported to be major phytoconstituents in stem and bark.

Seeds yield an alkaloid, a fatty oil, and a saponaceous glucoside. The alkaloid has properties identical to hypaphorine<sup>4</sup>.

Leaves and bark yield a poisonous alkaloid, erythrine, which acts on the nervous system with effects similar to the alkaloid cytosine. Bark, leaves and seeds yield saponin. Hydrocyanic acid is found in the leaves, stems, roots, and fruit. Phytochemical screening yielded eight spiroamine alkaloids and 3 carboxylated indole-3-alkylamines. Dried bark yields erythraline, hypaphorine, amino acids, organic acids, erythrinin, saponins<sup>5,6</sup>.

### Pharmacological actions

#### Anti Rheumatoid arthritis assay

The aqueous and ethanolic extracts of leaves of *Erythrina variegata* have been evaluated for the pharmacological studies by suspended in 4% v/v Tween 80. Both the extracts at the two different concentrations (256,500/kg body weight / oral once a day for 21 days) marked reduced swelling of the paw and bodyweight during secondary lesions. The EEEV is significantly higher. Finally WBC count. Together for the first time our studies have established the anti-arthritis potential of *Erythrina variegata* leaves extract<sup>7,8</sup>.

#### Anti microbial activity

The various extracts of *Erythrina variegata* were analysed against nine bacterial pathogens. Such as *salmonella typhi*, *pseudomonas putida*, *pseudomonas aeruginosa*, *E.coli*, *Klebsiella sp*, *Proteus sp*, *Serratia marcescens*, *Bacillus circulans*, *staphylococcus aureus* and *five fungal pathogens*, viz *Aspergillus niger*, *Aspergillus fumigatus*, *Sclerotium sp*, *Rhizopus sp*, *Candida sp*. All strains are more susceptible strains. Among all the solvents methanol showed promising anti microbial activity. These finally have potential for use in the treatment of diseases caused by these test organisms<sup>9</sup>.

#### Anti angiogenic activity

Three anthranoids were isolated from an alternaria Sp. Endophytic fungus is been evaluated for their anti angiogenic activity in a rat aortic sprouting assay, an *ex-vivo* model of angiogenesis of these three compounds alter solanoid was further characterised and found to show the activity in *ex-vivo*, *in-vitro* and *in-vivo* angiogenesis assays. The *in-vitro* model, the angiogenic effect of these was found in three main functions of endothelial cells, namely proliferation, tube formation and migration<sup>10,11</sup>.

#### Anti bacterial activity

The roots of *Erythrina variegata* were macerated with acetone. The chloroform soluble fraction of the residue was subjected to repeated silica gel column chromatography structures were isolated by elution with various solvents. Erycristagallin and orientanol B showed the highest anti MRSA activity<sup>12,13</sup>.

#### Anti convulsant activity

The chloroform extract of the root and bark of *Erythrina variegata* was been evaluated. The pentylenetetrazole and the maximal electric shock models were used for assessing the anticonvulsant effects of bark and root in mice and rats. Here, chloroform extract shows the onset of seizures compared with the control group in mice. Finally, it indicates that the chloroform root and bark extract of *Erythrina variegata* may be beneficial in both absence and tonic clonic seizures<sup>14,15</sup>.

### Diuretic activity

The chloroform and ethanol extract of *Erythrina variegata* was evaluated for the diuretic activity. The animals were divided into five groups deprived of food and water for 18 hours and the extracts were given to the animals (5 groups) according to their body weight the total volume of urine and the urine concentration of Na<sup>+</sup>, K<sup>+</sup> and Cl<sup>-</sup> and the Na<sup>+</sup> and K<sup>+</sup> were measured by  $\beta$  ame photometry<sup>16,17</sup>.

### CONCLUSION

In India, there are many tribals and other groups. These tribals are well aware of the usage of medicinal plants. Apart from that rural people also possess a good knowledge about the plants. In recent studies there was a much attention on these plants and their pharmacological studies so as to bring the known and unknown medicinal virtues especially of plant origin to lime light. In the present review article, we just had reviewed the literatures to collect the botanical, ethano botanical, phytochemical and pharmacological information on *Erythrina variegata*. Each part of the plant is used to treat various diseases in India. The different extracts of *Erythrina variegata* showed anti convulsant activity, anti bacterial activity, anti microbial activity, diuretic activity and etc. *Erythrina variegata* has many uses as a medicine yet its therapeutic efficacy has assessed majority in few cases.

### REFERENCES

1. Chopra RN, Nayar SL. Glossary of Indian medicinal plants, National institute of Science and Communication, Bangalore; 1996:111.
2. Ghani, medicinal plants of Bangladesh chemical constituents and uses, Asiatic society of Bangladesh, 1<sup>st</sup> edition 1988.
3. Suryawanshi H. P. Traditional uses, medicinal and phytopharmacological properties of *Erythrina indica* Lam: An overview. International journal of research in Ayurveda and pharmacy. 2011; 2(5): 1531-1533
4. Irfan Ali Khan, Atiya Khanum. Herbal medicines and uses Volume I, 1<sup>st</sup> edition. Ukaaz publications 2005: 34, 126, 139.
5. Sharma R, Medicinal plants of India, on encyclopaedia, Daya publishing House. Delhi 2003:99-100.
6. Tanaka H, Etoh H, Shimizu H, Makitha T, Talish V. Two new Isoflavonoids from *Erythrina variegata*. Planta medica. 2000:66 578-79.
7. Nadkarni K.M. The Indian material medical popular prakasha publishers Bombay. Volume 2000.
8. Suriyavanthana Muthu Krishnan Subha palanisamy. "Phytochemical screening and antimicrobial activity of *Erythrina variegata* world Journal of pharmacy and pharmaceutical sciences", 2014 volume: 3, Issue 6, 680-690.
9. Venkatesan D and Karrunakaran C.M. "Antimicrobial activity of selected Indian Medicinal plants" Journal of phytochemistry 2(2): 44-48 (2010).
10. He, Z. H, M.F, Ma S.C, butt P.P.H., Anti angiogenic effect of rhubarb and its anthraquinone derivatives. Journal of Ethanopharmacology 121, 313-317, 2009.
11. Strobel. G, Daisy. B, Castillo. U, Harper. Natural products from endophytic micro organisms, Journal of natural products 67, 257, 268. 2007.
12. Baitee, G.R and Neal, D. Vancomycin ototoxicity and nephrotoxicity. A review in medicinal toxicology -3, 376-386, 1988.
13. H. Tanaka, M. Sato. Anti bacterial activity of Isoflavonoids isolated from *Erythrina variegata* against Methicillin-resistant *Staphylococcus aureus*. 35, 494-498. 2002.
14. Chinchawade. A "International journal of pharmaceutical and clinical Research" 2013: 5(1) : 23-25.

15. Sander swas, sharvon SD, Epidemiology of Epilepsies J. Neuronal neuros psychiatry 1996: 61: 433-443.
16. Liu, I.X., Durham, D.G. and Richards, M.E., 2001. Vancomycin resistance reversal in enterococci by flavonoids. Journal of Pharmacy and Pharamacology., 53: 129-132.
17. Phunlap Pompenga et al. Antiangiogenetic effects of anthranoids from *Alternaria* sp., an endophytic fungus in a Thai medicinal plant *Erythrina variegata*. Phytomedicine. Volume 20, Issue 10, 15 July 2013, Pages 918–922

**Cite this article as:**

Lahari K, Divya M, Vidyavathi N, Kishore L, Poojitha M. A novel review on *Erythrina variegata*. Int. Res. J. Pharm. 2015; 6(4): 231-233 <http://dx.doi.org/10.7897/2230-8407.06451>

Source of support: Nil, Conflict of interest: None Declared