



## Review Article

### A REVIEW ON ANTHOCYANINS: A PROMISING ROLE ON PHYTOCHEMISTRY AND PHARMACOLOGY

Ashya Shaik <sup>1\*</sup>, Kishore Naidu Killari <sup>2</sup>, Jagadeesh Panda <sup>3</sup>

<sup>1</sup>Assistant Professor, Raghu College of Pharmacy, Dakamarri, Visakhapatnam, A.P, India

<sup>2</sup>Research scholar, AU College of Pharmaceutical Sciences, Andhra University, Visakhapatnam, A.P, India

<sup>3</sup>Principal, Raghu college of Pharmacy, Dakamarri, Visakhapatnam, A.P, India

\*Corresponding Author Email: shaikaasya@gmail.com

Article Received on: 03/12/17 Approved for publication: 22/01/18

**DOI: 10.7897/2230-8407.0911**

#### ABSTRACT


This paper reviews the literature on occurrence of different colors of anthocyanins, different types of anthocyanins containing food materials and their pharmacological actions. These anthocyanins are naturally available colored compounds from different flowers, fruits, and vegetables. Apart from various major pharmacological actions like anti-cancer, anti-inflammatory, decrease the neuronal death, these anthocyanins are acts as natural colorants for food and beverages. There is no study reveal that toxic effect of anthocyanins on human health. However, future research is going on pharmacological actions and uses of anthocyanins.

**Key Words:** Anthocyanins, Transformation, Pharmacological effects

#### INTRODUCTION

Pigmentation is an attractive feature of various fruits, flowers and other main substances. Anthocyanins (Anthos = flower and kianos = blue) are one of the responsible for red-blue coloration, these are the most important pigments of the vascular plants, they are harmless and easy dissolve in aqueous media, so these are known as natural water-soluble colorants. These anthocyanins are most utilized vegetable colorants. In the international food and beverage industry's anthocyanins preferred natural food colors like orange, red, pink and purple to blue<sup>1</sup>. They were used by Americans, Indians, Europeans, Japanese and the Chinese. These water-soluble anthocyanins are low stable than carotenoids. They belong to a molecule called flavonoids synthesized via the phenylpropanoid pathway they are odorless and nearly flavorless, its taste as a moderately astringent sensation. They are extracted from wide varieties of fruits, vegetables and other food materials like grapes, berries, red cabbage, apples, radishes, tulips, roses and orchids, olive oil, honey, teas, cocoa, banana, asparagus, pea, fennel, pear and potato. Recent experimental studies declared, compounds of colored fruits, berries and vegetables, which may scavenge free radicals very strongly (antioxidant effect) and prevent distinct chronic diseases like protection against liver disorders, hypertension, enhance memory, progress eyesight, anti-inflammatory and antimicrobial activities, inhibition of mutations caused by mutagens from baked food, and suppression against tumors. Other than these the consumption of anthocyanins may play a significant character in preventing lifestyle-related diseases such as cancer, hyperglycemia, and cardiovascular and neurological disorder. Anthocyanins are derived from anthocyanidins by adding sugars

#### BASIC STRUCTURE OF ANTHOCYANIN



**Figure 1: Basic structure of anthocyanins and its substituents<sup>2</sup>**

Sugar free anthocyanins are anthocyanidins. The anthocyanins are classified into 2 types they are sugar- free anthocyanidin aglycons and the anthocyanin glycosides. Till now 500 types of anthocyanins are reported. Examples of sugar free anthocyanidin aglycons like: Delphinidin (blue-red), Pelargonidin (orange) and Cyanidin (orange-red). These sugar free anthocyanidin aglycons are the building blocks which are produce anthocyanins by reduction, dehydroxylation and glycosylation within the plant.

#### BIOSYNTHESIS OF ANTHOCYANINS

Anthocyanins are formed from two different parts of chemical raw materials which is present in cell: From this one part involves the production of the phenylalanine by shikimate pathway. Other part involves produce three molecules of malonyl-coA, a C3 unit

from a C2 unit (acetyl-coA)<sup>3</sup>. These two steps are combining to form an intermediate chalcone-like compound in the presence of chalcone synthase enzyme by using polyketide folding mechanism that is commonly found in plants. The resulting product chalcone is isomerized as prototype naringenin pigment in the presence of chalcone isomerase. The formed Naringenin product is subsequently oxidized in the presence of flavanone hydroxylase, flavonoid 3' hydroxylase and flavonoid 3' 5'-hydroxylase enzymes. These resulting products are further reduced in presence of dihydroflavonol 4-reductase then forms a colorless leucoanthocyanidins<sup>4</sup>. Leucoanthocyanidins are precursors of the next enzyme, anthocyanidin synthase or leucoanthocyanidin dioxygenase. Flavan-3-ols, the products of leucoanthocyanidin reductase (LAR), have been recently shown to be their true substrates then unstable anthocyanidins are formed. These unstable anthocyanidin further coupled to sugar molecules by enzymes such as UDP-3-O-glucosyltransferase to yield the final relatively stable anthocyanins<sup>5</sup>.

## FUNCTIONS OF ANTHOCYANINS

### Anthocyanins used as food additive

Mainly color plays a crucial role in food industries. Acceptability of food is mostly based on appearance of food. So many industries use synthetic food colorants to improve the elegance of food. These synthetic colorants show many side effects compared to natural colorants which are synthesized from anthocyanins. They show many beneficial effects without adverse effects. Anthocyanins involve a distinct group of intensely colored pigments responsible for the orange, red, purple and blue colors. These colors extracted from many leaves fruits, vegetables, flowers, roots<sup>6</sup> Cyanidin, Delphinidin and Pergolidin are the anthocyanins which are responsible for the pigmentation. They are extracted from strawberry, radish, potato, red grapes, cranberry<sup>7</sup> blueberries<sup>8</sup>, black chokeberries<sup>9</sup>, elderberries, black currents<sup>10</sup>, purple corn<sup>11</sup>.

### Free radical scavenging activity of anthocyanins

Free radical is any atom or molecule that contains one or more unpaired electrons. Anthocyanins scavenge free radicals shows an antioxidant activity by increased uric acid. Low-wavelength electromagnetic radiation (eg. gamma rays) can split water in the body to generate hydroxyl radical. Apart from this another radical are oxygen radical ( $O_2$ ) and nitric oxide. These super oxide and nitric oxide has beneficial effects in low concentration. radicals can generate more toxic chemicals in traces amounts. These two free radicals combine to form peroxy radical, which damage the proteins and react with the iron copper<sup>12</sup>. Hydrogen peroxide radical also form during the stress condition. These are highly reactive mainly in nucleus and damaging the membrane of DNA proteins carbohydrates, lipids and target all kind of cells in body. They are produced from vital metabolic activity in body and some of the external source also responsible for the free radical formation X-rays, ozone, cigarette smoking, air pollutants, industrial chemicals, mitochondria, xanthine oxidase, peroxisome, inflammation, phagocytosis, exercise, ischemic injury, certain drug and pesticide. These free radicals may cause different diseases in body like cancer, ischemia disease. Anthocyanins rich food contains a cyanidin, delphinidin, malvidin, peonidin these all are pays a crucial role in scavenge free radicals and reduces the lipid peroxidation and DNA damage<sup>13</sup>. For example, eggplant shows anti-oxidant activity by scavenge the oxygen free radical<sup>14</sup> black berries having the highest free radical scavenging capacity against super oxide radical, hydrogen peroxide radical and nitric oxide radical<sup>15</sup> and red cabbage containing anthocyanins are protect from oxidative stress<sup>16</sup>. The anthocyanidin pelargonidin

protect the amino acid tyrosine by highly reactive peroxy radical free radical, anthocyanidin like delphinidin called nasunin, which destroy the dangerous hydroxyl radical<sup>17</sup>. These anthocyanidin shows a four times powerful antioxidant activity than vitamin E<sup>18</sup>. Philippines Medusa, Siberian Ginseng, arabidopsis, red onions, chagalopoli, asargu, red cabbage broccoli tea, red orange saffron black carrot, Ceylon goose berry, palm fennel, straw berry, black soybean, sweet potato, Chinese bay berry, myrtle berry, orchids, black rice, sweet cherry, sour cherry, pomegranate, peach, radish, goose berry, black current, rose petals, black raspberry, tomato, elderberry, bilberry, bog berry, wheat, grapes, corn. These are the anthocyanins rich food which will discuss in below Table 2.

### Antihyperglycemic activity of anthocyanins

High fat diet is the one of the risk factor of type- ii diabetes in this insulin resistance results hyperglycemia occur. Anthocyanin like cyaniding, Delphinidin, pionidin, pelargonidin effectively treat the hyperglycemia by decreasing the tumor necrosis factor (TNF)- $\alpha$  mRNA levels and decreasing oxidative stress protect pancreatic  $\beta$  cells<sup>19</sup> and also it decreases the glucose production by increases the AMPK, increase the gucose absorption and transport then lower risk of diabetes<sup>20</sup>. For example anthocyanin which extract from Purple corn, cornus fruits (cherry), a rich source of anthocyanins, chockberry, litchi, cranberry shows better hypoglycemic activity. Apart from anti hyperglycemic activity these anthocyanins prevent the symptom which are associate with diabetes like diabetic retinopathy<sup>21</sup>. One research article proved that anthocyanins which are present in back soybean seed upper layer acts against type-ii diabetes through regulation of glucose transporter 4 and inhibit pancreatic apoptosis<sup>22</sup>.

### Effect of anthocyanin on Cardio vascular system

Oxidative stress is one of the risk factor for cardiovascular disease. Anthocyanins play a vital role in protection against oxidative stress. The anthocyanin which is extracted from elder berries acts on endothelial cells shows protective effects<sup>23</sup>. Delphinidin, which is extracted from red wine having vasodilation activity<sup>24</sup>. We already know hyperlipidemia is a dangerous risk factor for cardiovascular diseases, cyaniding rich content containing black current could be decrease the fatty acid levels<sup>25</sup>. A strong evidence is there anthocyanin rich content containing grape juice have powerful antioxidant activity protect against heart attack through increase the capillary permeability and strength, and to inhibit platelet formation and increase nitric oxide (NO) production the vasodilatation occurs<sup>26</sup>. These anthocyanins reduce active against lipid peroxidation then protect the cell membrane finally prevent the blood vessel damage and DNA dmgage<sup>27</sup>. The Anthocyanins are preventing the plaque formation in arteries. For example, delphinidin is an anthocyanidin, it relaxes the blood vessels and it reduces the cardiovascular mortality". During the ischemia-reperfusion free radicals are form that result in white blood cell adhesion to micro capillary walls, then capillary wall permeability increases, blood flow decreases, finally causes permanent capillary damage these anthocyanins also helped in reduces the capillary damage<sup>28</sup>. Anthocyanins scavenge the free radicals then inhibit the reactive oxygen species finally inhibit the inflammation related cardiovascular diseases.

### Effect of anthocyanin on nervous system

Oxidative stress is a major factor for many neurodegenerative disorders. Anthocyanins give potent relief from this oxidative damage, for example Delphinidin is an anthocyanidin which protects lipids in brain tissue from deterioration. Pelargonidin may help protect against neurological diseases by preventing

Peroxynitrite free radical. Straw berry extract contains anthocyanidins like pelargonidin and delphinidine which shows a neuroprotective activity

#### Anti-inflammatory activity of anthocyanins

During the inflammation condition increased production of nitric oxide free radicals and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) in presence of Cyclooxygenase (COX), mainly COX-2 is a gene that is highly inducible by inflammatory stimuli. Mainly anthocyanins show anti-inflammation activity by inhibits COX pathway very effectively. For example *Punica granatum* shows anti-inflammatory activity, cyanidin glycosides containing cherries and berries also shows anti-inflammatory activity, Siberian ginseng containing anthocyanidin like cyaniding shows anti-inflammatory activity, back soybean and ipomea containing anthocyanidin like Cyanidin, Delphinidin, Petunidin, and Pelargonidin, Jabuticaba containing cyanidin and delphinidin and Ceylon gooseberry containing Cyanidin, Peonidin, Malvidin, Petunidin, Delphinidin and Pelargonidin how potent anti-inflammatory effect<sup>29</sup>.

#### Effect of anthocyanins on eye

The anthocyanins may also increase eyesight and treat eye infections. A recent study found that black currant and maqui berry containing anthocyanidins used to improve eyesight<sup>30</sup>.

#### Effect of anthocyanins on cancer

Anthocyanins may have capability to treat the tumorigenesis and cell proliferation<sup>31-33</sup>. It inhibits the carcinogenesis through many multiple mechanisms are antioxidant activity and potent COX enzyme inhibition<sup>34</sup>. Anthocyanin act against tumors by blocking MAP kinase pathway. For example, the anthocyanins which shows anticancer activity are Siberian Ginseng, Black carrot, Palm, Ipomea, Honey berry<sup>35,36</sup>.

#### Effect of anthocyanins on ulcers

The anthocyanin like Bilberry, used for to treat ulcers by the increasing the production of stomach mucus and protect the stomach from injury. Elderberry extract reduced stress-induced ulcers<sup>37</sup>.

#### Other functions of anthocyanins

Apart from above functions anthocyanins protects from microbial infections, diarrhea, anti-fungal, immunomodulatory action, analgesic, Hypertension, Dysentery<sup>38</sup>. List of anthocyanins described in Table 2.

Table 1: Selected Anthocyanidins and their Substituents

Anthocyanidin	R <sub>3</sub> '	R <sub>4</sub> '	R <sub>5</sub> '	R <sub>3</sub> '	R <sub>5</sub> '	R <sub>6</sub> '	R <sub>7</sub> '
Aurantidin	-H	-OH	-H	OH	-OH	-OH	-OH
Cyanidin	-OH	-OH	-H	OH	-OH	-H	-OH
Delphinidin	-OH	-OH	-OH	OH	-OH	-H	-OH
Europinidin	-OCH <sub>3</sub>	-OH	-OH	OH	-OCH <sub>3</sub>	-H	-OH
Pelargninidin	-H	-OH	-H	OH	-OH	-H	-OH
Malvidin	-OCH <sub>3</sub>	-OH	-OCH <sub>3</sub>	OH	-OH	-H	-OH
Peonidin	-OCH <sub>3</sub>	-OH	-H	OH	-OH	-H	-OH
Petunidin	-OH	-OH	-OCH <sub>3</sub>	OH	-OH	-H	-OH
Rosinidin	-OCH <sub>3</sub>	-OH	-H	-OH	-OH	-H	-OCH <sub>3</sub>

Table 2: List of Anthocyanin Containing Food Materials and their Pharmacological Role

Source	Anthocyanins	Function
<i>Acalypha hispida</i> <sup>39</sup> (Philippines Medusa)	Cyanidin	Anti-fungal and Anti-oxidant activity
<i>Allium cepa</i> <sup>40</sup> (Red onions)	Cyanidin and Peonidin	Anti-oxidant activity
<i>Arabidopsis thaliana</i> <sup>41</sup> (Arabidopsis)	Cyanidin	Antioxidative activity
<i>Ardisia compressa</i> K <sup>42</sup> (Chagalapoli)	Malvidin, petunidin and Delphinidin	Anti-oxidant activity
<i>Aristotelia chilensis</i> <sup>43</sup> (Maqui berry)	Delphinidin	Treat eye diseases
<i>Aronia melanocarpa</i> <sup>44</sup> (Choke berries)	Cyanidin	Cardio-protective, Antidiabetic and Immunoregulatory effects
<i>Asparagus densiflorus</i> <sup>45</sup> (Asparagus)	Cyanidin and Peonidin	Anti-oxidant activity
<i>Begonia semperflorens</i> <sup>46</sup> (Wax begonia)		Photo-protective capacity
<i>Brassica oleracea</i> <sup>47</sup> (Red cabbage)	Cyanidin	Anti-oxidant activity
Broccoli <sup>48</sup>	Cyanidin	Antioxidant activity
<i>Camellia hongkongensis</i> <sup>49</sup>	Cyanidin and Delphinidin	Antioxidant activity
<i>Camellia sinensis</i> <sup>50</sup> (Teas)	Delphinidin and Cyanidin	Anti-oxidant activity
<i>Citrus sinensis</i> <sup>51</sup> (Blood orange)	Cyanidin	Antioxidant activity
<i>Crataegus laevigata</i> <sup>52</sup> (Haw thorne)	Cyanidin	Anti-oxidant, Anti-microbial and Protect against cardiovascular diseases

<i>Crocus sativus</i> <sup>53</sup> (saffron tepals)	Cyanidin, Pelargonidin, Delphinidin and Petunidin	Anti-oxidant activity
<i>Daucus carota</i> <sup>54</sup> (Black carrot)	Cyanidin	Anti-cancer and Anti-oxidant activity
<i>Dovyalis hebecarpa</i> <sup>55</sup> (Ceylon gooseberry)	Delphinidin and cyaniding	Antioxidant activity
<i>Erythrina crista-galli</i> L <sup>56</sup> (cockspur coral tree)	Cyanidin	Astringent, Narcotic, and Analgesic for wound healing
<i>Euterpe oleracea</i> <sup>57</sup> (Palm)	Cyanidin, pelargonidin and peonidin	Anticancer and Antioxidant effects
<i>Foeniculum vulgare</i> <sup>58</sup> (Fennel)	Cyanidin	Anti-oxidant and Anti-microbial activity
<i>Fragaria ananassa</i> <sup>59</sup> (Strawberry)	Cyanidin and Pelargonidin	Neuroprotectant activity
<i>Fragaria chiloensis</i> <sup>60</sup> (Chilean strawberry)	Cyanidin and Pelargonidin	Anti-oxidant activity
<i>Garcinia indica</i> <sup>61</sup> (Kokum)	Cyanidin	Treat Dysentery, Tumours, Heart complaints, Stomach acidity and Liver disorders
<i>Glycine max</i> L <sup>62</sup> (Black soybean)	Cyanidin, Delphinidin, Petunidin, and Pelargonidin	Antioxidant activity, protect from Ischemia, Reperfusion, Heart injury, Anti-inflammation and Wound healing
<i>Gynura bicolor</i> <sup>63</sup>	Cyanidin	Antioxidant and Anti-cancer activity
<i>Hemigraphis colorata</i> <sup>64</sup>	Cyanidin	Anti-bacterial activity
<i>Hibiscus sabdariffa</i> <sup>65</sup> (Roselle calyces)	Delphinidin and Cyanidin	Used in the food, pharmaceutical and cosmetic industries
<i>Ipomoea batatas</i> <sup>66</sup> (Sweet potato).	Peonidin and Cyanidin	Anti-oxidant activity
<i>Litchi chinensis</i> Sonn <sup>67</sup> (litchi)	Cyanidin, Malvidin	Anti-oxidant, Anti-diabetic, Antimicrobial, Anti ulcers, Anti-obesity, having analgesic action, Antihyperglycemic, Antihyperlipidemic, Antiplatelet, Antiviral and Hepatoprotective activity
<i>Lonicera Caerulea</i> <sup>68</sup> (Honey berry)	Cyanidin	Chemopreventive, antimicrobial, antiadherence and antioxidant effects
<i>Lonicera japonica</i> <sup>69</sup> (Honeysuckle)	Cyanidin	Anti-oxidant activity
<i>Matthiola fruticulosa</i> <sup>70</sup> (Sad stock)	Cyanidin	Antioxidant activity
<i>Matthiola tricuspidata</i> <sup>71</sup> (Three-horned stock)	Cyanidin	Antioxidant activity
<i>Morus australis</i> Poir <sup>72,73</sup> (Mulberry)	Cyanidin and Pelargonidin	Anti-fever diuretics, Liver protection, Anti-hypertensive, Anti-obesity And protects from Cardiovascular diseases
<i>Musa paradisiaca</i> <sup>73,74</sup> (Banana)	Cyanidin, Delphinidin, Pelargonidin, Peonidin, Petunidin and Malvidin	Anti-ulcer
<i>Myrciaria cauliflora</i> <sup>75</sup> (Jaboticaba)	Cyanidin and Delphinidin	Treat hemoptysis, Asthma, Diarrhea and Gargled for chronic inflammation of the tonsils
<i>Myrica rubra</i> Sieb <sup>76</sup> (Chinese bayberry)	Cyanidin	Anti-oxidant activity
<i>Myrtus communis</i> <sup>77</sup> (Myrtle berries)	Delphinidin, Petunidin, Malvidin, Peonidin and Cyanidin	Antioxidative activity
<i>Neomitrantes</i> <sup>78</sup> <i>Obscura</i>	Cyanidin, Delphinidin, Peonidin and Pelargonidin	Antioxidant activity
<i>Ocimum basilicum</i> <sup>79</sup> (Basil)	Cyanidin	Antioxidant activity and Anti- Parkinson's disease
<i>Orchis</i> <sup>80</sup> (Orchids)	Cyanidin	Anti-oxidant activity
<i>Oryza sativa</i> <sup>81</sup> (Black rice)	Cyanidin and Peonidin	Anti-oxidant activity
<i>Paeonia suffruticosa</i> <sup>82,83</sup> (Peony)	Pelargonidin, Cyanidin and Peonidin.	Acaricidal activity
<i>Phaseolus vulgaris</i> <sup>84</sup> (Black bean)	Delphinidine, Malvidine, Petunidin	Anti-cancer
<i>Pisum sativum</i> <sup>85</sup> (peas)	Cyanidin	Anti-oxidant activity
<i>Plumeria rubra</i> <sup>86</sup> (Red frangipani)	Cyanidin	Antioxidant and hypolipidemic activity
<i>Prunus avium</i> <sup>87</sup> (Sweet cherries)	Cyanidin	Anti-oxidant activity
<i>Prunus cerasus</i> var. <i>Marasca</i> <sup>88</sup> (Sour cherry Marasca)	Cyanidin	Anti-oxidant activity
<i>Prunus mahaleb</i> <sup>89</sup> (Rock cherry)	Cyanidin, Peonidin and Pelargonidin	Natural pigment for food industry

<i>Prunus persica</i> <sup>90</sup> (peach)	Peonidin and Cyanidin	Anti-oxidant and Anti-ulcers
<i>Punica granatum</i> <sup>91,92</sup> (Pomogante)	Delphinidin, Cyanidin and Pelargonidin	Protection against Cardiovascular diseases, Antioxidant activity and Anti- cancer
<i>Pyrus communis</i> <sup>93</sup> (Pear)	Cyanidin and Peonidin	Anti-cancer activity
<i>Queen Garnet</i> <sup>94</sup> (plum)	Cyanidin	Antithrombotic and Anti-oxidant activity
<i>Raphanus sativus</i> <sup>95</sup> (Radish)	Cyanidin and pelargonidin	Anti-oxidant activity
<i>Rhodomyrtus tomentosa</i> <sup>96</sup> (Ceylon hill gooseberry)	Cyanidin, Peonidin, Malvidin, Petunidin, Delphinidin and Pelargonidin	Antioxidant, Anticancer Anti-inflammatory, Anti artery- atherosclerosis, Anti-hypertensive and Antibacterial activities
<i>Ribes nigrum</i> <sup>97</sup> (Blackcurrant)	Delphinidin and Cyanidin	Anti-oxidant, Anti-coagulant, Anti-hypertensive, Anti-inflammatory, Anti-microbial, Anti-thrombotic and Anti-viral activities
<i>Rosa hybrida</i> <sup>98</sup> (Rose)	Cyanidin, Pelargonin and Peonidin	Anti-inflammatory and Anti-oxidant activities
<i>Rubus fruticosus</i> <sup>99,100</sup> (Black berry)	Cyanidin, Pelargonidin, and Peonidin	Anti-oxidant property, Protection against endothelial dysfunction, protect against vascular dysfunction, and Anti-cancer
<i>Solanum lycopersicum. L</i> <sup>101</sup> (Tomato)	Aurantininidin, Cyanidin, Delphinidin, Pelargolidin, Malvidin, Peonidin, Petunidin and Rosinidin	Anti-oxidant activity and protect against cardiac injury
<i>Solanum melongena</i> <sup>102</sup> (Eggplant)	Delphinidin	Anti-oxidant activity, anti-diabetic activity
<i>Solanum tuberosum</i> <sup>103</sup> (Potato)	Pelargonidin, Petunidin, Cyanidin, and Malvidin	Protect against heart disease and Anti- cancer
<i>Syzygium cumini</i> <sup>104</sup> (Jamun)	Malvidin	Anti-oxidant, Antifungal, Anti-diabetic and Anti-inflammatory activities
<i>Triticum aestivum</i> <sup>105</sup> (Canada's spring wheat)	Cyanidin, Delphinidin and Peonidin	Antioxidant activity
<i>Vaccinium myrtillus</i> <sup>106</sup> (Bil berry)	Pelargolidin, Cyanidini, Peonidin, Delphinidin and Malvidin	Antioxidant, Antihyperglycemia, Antihyperlipidemia, Anti-Cancer Improving vision, antiinflammatory, Dementia, Cardio Vascular disease, and other age-related diseases.
<i>Vaccinium uliginosum</i> <sup>107</sup> (Bog berry)	Delphinidin, Cyanidin, Petunidin, Peonidin, and Malvidin	Antioxidant, Anti-inflammatory and Anti-cancer
<i>Vaccinum corymbosum</i> <sup>108</sup> (Blue berry)	Cyanidin, Delphinidin, Petunidin, Peonidin and Malvidin	Anticancer, vascular diseases and Neurodegenerative diseases
<i>Vaccinum macrocarpn</i> <sup>109</sup> (Cran berry)	Cyanidin, Delphinidin and peonidin	Treat cancer, inflammation, dyslipidemia, hyperglycemia and oxidative stress, UTI, cardiac Diseases
<i>Vitis vinifera</i> <sup>110</sup> (Grapes)	Malvidin, Cyanidin and Peonidin	Anti-oxidant activity
<i>Zea mays</i> (corn)	Cyanidin	Antioxidant activity

## SUMMARY

The improvement of living conditions has contributed to the increased longevity of people worldwide. As a result, population aging is no longer a phenomenon restricted to developed countries.

In recent years, studies have begun to investigate the specific properties of isolated anthocyanin pigments. However, many still study the health effects of anthocyanins from fruit extracts where anthocyanins are present in combination with other compounds. In fact, some reports suggest that anthocyanin activity is actually potentiated when delivered in mixtures, as opposed to isolate.

## ACKNOWLEDGEMENT

The corresponding author is obliged to Dr Killari Kishore Naidu, AU College of Pharmaceutical Sciences, Andhra University for support the necessary facilities to complete the present review.

## REFERENCES

1. Wills JM, Schmidt DB, Pillo-Blocka F, Cairns G. Exploring global consumer attitudes toward nutrition information on food labels. *Nutrition Reviews* 2009; 67(1):102-6.
2. Kong JM, Chia LS, Goh NK, Chia TF, Brouillard R. Analysis and biological activities of anthocyanins. *Phytochemistry* 2003; 64(5): 923-33.
3. Klaus M. Herrmann, Lisa M. Weaver. The shikimate pathway. *Annual review of plant physiology and plant molecular biology* 1999; 50: 473-503
4. Nakajima JI, Tanaka Y, Yamazaki M, Saito K. Reaction mechanism from leucoanthocyanidin to anthocyanidin 3-glucoside, a key reaction for coloring in anthocyanin biosynthesis. *Journal of Biological Chemistry* 2001; 276(28): 25797-803.
5. Kovinich N, Saleem A, Arnason JT, Miki B. Functional characterization of a UDP-glucose: flavonoid 3-O-glucosyltransferase from the seed coat of black soybean

- (Glycine max (L.) Merr.). *Phytochemistry*. 2010; 71(11): 1253-63.
6. Bridle P, Timberlake CF. Anthocyanins as natural food colours—selected aspects. *Food chemistry* 1997;58(1): 103-9.
  7. J.Côté, S.Caillet, D.Dussault, J-F.Sylvain, M.Lacroix. Effect of juice processing on cranberry antibacterial properties. *Food Research International* 2011; 44 (9): 2922-2929.
  8. Francis FJ. Blueberries as a colorant ingredient in food products. *Journal of Food Science* 1985;50(3):754-6.
  9. Giusti MM, Wrolstad RE. Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal* 2003;14(3): 217-25.
  10. Jordi Giné Bordonaba and Leon A. Terry. Biochemical Profiling and Chemometric Analysis of Seventeen UK-Grown Black Currant Cultivars. *Journal of agricultural and food chemistry* 2008; 56 (16): 7422–7430.
  11. Nakatani N, Fukuda H, Fuwa H. Major anthocyanin of Bolivian purple corn (*Zea mays* L.). *Agricultural and Biological Chemistry* 1979;43(2): 389-91.
  12. Halliwell B. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence. *The lancet*. 1994;344 (8924): 721-4.
  13. Ramirez-Tortosa C, Andersen ØM, Gardner PT, Morrice PC, Wood SG, Duthie SJ, Collins AR, Duthie GG. Anthocyanin-rich extract decreases indices of lipid peroxidation and DNA damage in vitamin E-depleted rats. *Free Radical Biology and Medicine*. 2001;31 (9): 1033-7.
  14. Sadilova E, Stintzing FC, Carle R. Anthocyanins, colour and antioxidant properties of eggplant (*Solanum melongena* L.) and violet pepper (*Capsicum annuum* L.) peel extracts. *Zeitschrift für Naturforschung C* 2006; 61(7-8): 527-35.
  15. Wang SY, Jiao H. Scavenging capacity of berry crops on superoxide radicals, hydrogen peroxide, hydroxyl radicals, and singlet oxygen. *Journal of Agricultural and Food Chemistry* 2000;48(11): 5677-84.
  16. Igarashi K, Kimura Y, Takenaka A. Preventive effects of dietary cabbage acylated anthocyanins on paraquat-induced oxidative stress in rats. *Bioscience, biotechnology, and biochemistry*. 2000; 64(8): 1600-7.
  17. Noda Y, Kneyuki T, Igarashi K, Mori A, Packer L. Antioxidant activity of nasunin, an anthocyanin in eggplant peels. *Toxicology* 2000;148(2): 119-23.
  18. Tsuda T, Horio F, Kitoh J, Osawa T. Protective effects of dietary Cyanidin 3-O-β-d-Glucoside on liver ischemia-reperfusion injury in rats. *Archives of Biochemistry and Biophysics*. 1999; 368(2): 361-6.
  19. Tsuda T, Horio F, Uchida K, Aoki H, Osawa T. Dietary cyanidin 3-O-β-D-glucoside-rich purple corn color prevents obesity and ameliorates hyperglycemia in mice. *The Journal of nutrition* 2003; 133(7):2125-30.
  20. Kruger MJ, Davies N, Myburgh KH, Lecour S. Proanthocyanidins, anthocyanins and cardiovascular diseases. *Food Research International*. 2014; 59: 41-52.
  21. Perossini M. Diabetic and hypertensive retinopathy therapy with *Vaccinium myrtillus* anthocyanosides (Tegens). *Annali Di Ottalmologia E Clinica Oculistica* 1987; 113: 1173.
  22. Nizamudinova IT, Jin YC, Chung JI, Shin SC, Lee SJ, Seo HG, Lee JH, Chang KC, Kim HJ. The anti-diabetic effect of anthocyanins in streptozotocin-induced diabetic rats through glucose transporter 4 regulation and prevention of insulin resistance and pancreatic apoptosis. *Molecular nutrition & food research* 2009;53(11): 1419-29.
  23. Youdim KA, Martin A, Joseph JA. Incorporation of the elderberry anthocyanins by endothelial cells increases protection against oxidative stress. *Free Radical Biology and Medicine* 2000;29(1): 51-60.
  24. Andriambeloston E, Magnier C, Haan-Archipoff G, Lobstein A, Anton R, Beretz A, Stoclet JC, Andriantsitohaina R. Natural dietary polyphenolic compounds cause endothelium-dependent vasorelaxation in rat thoracic aorta. *The Journal of nutrition*. 1998; 128(12): 2324-33.
  25. Frank J, Kamal-Eldin A, Lundh T, Määttä K, Törrönen R, Vessby B. Effects of dietary anthocyanins on tocopherols and lipids in rats. *Journal of agricultural and food chemistry* 2002; 50(25): 7226-30.
  26. Folts JD. Antithrombotic potential of grape juice and red wine for preventing heart attacks. *Journal of Pharmaceutical Biology* 1998; 36: 21-7.
  27. Ramirez-Tortosa C, Andersen ØM, Gardner PT, Morrice PC, Wood SG, Duthie SJ, Collins AR, Duthie GG. Anthocyanin-rich extract decreases indices of lipid peroxidation and DNA damage in vitamin E-depleted rats. *Free Radical Biology and Medicine* 2003;31(9): 1033-7.
  28. Bertuglia S, Malandrino S, Colantuoni A. Effect of *Vaccinium myrtillus* anthocyanosides on ischaemia reperfusion injury in hamster cheek pouch microcirculation. *Pharmacological research*. 1995; 31(3): 183-7.
  29. Seeram NP, Adams LS, Zhang Y, Lee R, Sand D, Scheuller HS, Heber D. Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells in vitro. *Journal of agricultural and food chemistry*. 2006; 54(25): 9329-39.
  30. Nakaishi H, Matsumoto H, Tominaga S, Hirayama M. Effects of black currant anthocyanoside intake on dark adaptation and VDT work-induced transient refractive alteration in healthy humans. *Alternative Medicine Review* 2000;5(6):553-62.
  31. Stoner GD, Sardo C, Apseloff G, Mullet D, Wargo W, Pound V *et al*. Pharmacokinetics of Anthocyanins and Ellagic Acid in Healthy Volunteers Fed Freeze-Dried Black Raspberries Daily for 7 Days. *The Journal of Clinical Pharmacology*. 2005; 45(10): 1153-64.
  32. Kang SY, Seeram NP, Nair MG, Bourquin LD. Tart cherry anthocyanins inhibit tumor development in Apc Min mice and reduce proliferation of human colon cancer cells. *Cancer Letters - Journal* ;194(1): 13-9.
  33. Koide T, Hashimoto Y, Kamei H, Kojima T, Hasegawa M, Terabe K. Antitumor effect of anthocyanin fractions extracted from red soybeans and red beans in vitro and in vivo. *Cancer biotherapy & radiopharmaceuticals*. 1997;12 (4): 277-80.
  34. Meiers S, Kemény M, Weyand U, Gastpar R, von Angerer E, Marko D. The anthocyanidins cyanidin and delphinidin are potent inhibitors of the epidermal growth-factor receptor. *Journal of agricultural and food chemistry* 2001; 49(2): 958-62.
  35. Hou DX, Kai K, Li JJ, Lin S, Terahara N, Wakamatsu M, Fujii M *et al*. Anthocyanidins inhibit activator protein 1 activity and cell transformation: structure–activity relationship and molecular mechanisms. *Carcinogenesis*. 2004; 25(1): 29-36.
  36. Bomser J, Madhavi DL, Singletary K, Smith MA. In vitro anticancer activity of fruit extracts from *Vaccinium* species. *Planta medica* 1996; 62(03): 212-6.
  37. Kandil FE, Song L, Pezzuto JM, Marley K, Seigler DS, Smith MA. Isolation of oligomeric proanthocyanidins from flavonoid-producing cell cultures. *In Vitro Cellular & Developmental Biology-Plant*. 2000;36(6): 492-500.

38. RobertVeberic, JernejaJakopic, FranciStampar, Valentina, Schmitzer. European elderberry (*Sambucus nigra* L.) rich in sugars, organic acids, anthocyanins and selected polyphenols. Food Chemistry 2009; 114 (2): 511-515
39. Rice-evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. Free radical research 1995;22(4): 375-83.
40. Ejechi BO, Souzaey JA. Inhibition of biodeterioration of yam tuber *Dioscorea rotundata* Poir in storage with phenolic extract of *Acalypha hispida* Burm. F leaves. Journal of Stored Products Research. 1999;35(2): 127-34.
41. FOSSEN T, ANDERSEN ØM, ØVSTEDAL DO, PEDERSEN AT, RAKNES Å. Characteristic anthocyanin pattern from onions and other *Allium* spp. Journal of food science. 1996; 61(4): 703-6.
42. Ghassami F, Alavi A, Mirkheshti N, Ansari L. Study the therapeutic effect of Colchicine on delayed pulmonary injury of sulfur mustard in animal model. Trauma Monthly. 2008; 20(02): 125-32.
43. Joaquín-Cruz E, Dueñas M, García-Cruz L, Salinas-Moreno Y, Santos-Buelga C, García-Salinas C. Anthocyanin and phenolic characterization, chemical composition and antioxidant activity of chagalapoli (*Ardisia compressa* K.) fruit: A tropical source of natural pigments. Food Research International. 2015; 70: 151-7.
44. Tanaka J, Kadekaru T, Ogawa K, Hito S, Shimoda H, Hara H. Maqui berry (*Aristotelia chilensis*) and the constituent delphinidin glycoside inhibit photoreceptor cell death induced by visible light. Food chemistry. 2015;139(1): 129-37.
45. Kulling SE, Rawel HM. Chokeberry (*Aronia melanocarpa*)—A review on the characteristic components and potential health effects. Planta medica 2008;74(13): 1625-34.
46. Wann EV, Thompson AE. Anthocyanin pigments in asparagus. Proceedings of the Society for Horticultural Science 1965;87: 270-273.
47. Pietrini F, Iannelli MA, Massacci A. Anthocyanin accumulation in the illuminated surface of maize leaves enhances protection from photo-inhibitory risks at low temperature, without further limitation to photosynthesis. Plant, Cell & Environment. 2002; 25(10): 1251-9.
48. Cevallos-Casals BA, Byrne D, Okie WR, Cisneros-Zevallos L. Selecting new peach and plum genotypes rich in phenolic compounds and enhanced functional properties. Food Chemistry. 2006;96 (2): 273-80.
49. Li JB, Hashimoto F, Shimizu K, Sakata Y. A new acylated anthocyanin from the red flowers of *Camellia hongkongensis* and characterization of anthocyanins in the section *Camellia* species. Journal of integrative plant biology. 2009; 51(6): 545-52.
50. Nanjo F, Mori M, Goto K, HARA Y. Radical scavenging activity of tea catechins and their related compounds. Bioscience, biotechnology, and biochemistry. 1999; 63(9): 1621-3.
51. Maccarone E, Rapisarda P, Fanella F, Arena E, Mondello L. Cyanidin-3-(6-malonyl)- $\beta$ -glucoside. One of the major anthocyanins in blood orange juice. Italian journal of food science. 1998; 10(4): 367-72.
52. Saeb H, Khayyat M, Zarezadeh A, Moradinezhad F, Samadzadeh A, Safae M. Effects of NaCl Stress on Seed Germination Attributes of Periwinkle (*Catharanthus Roseus* L.) and Corn Poppy (*Papaver Rhoëas* L.) Plants. Plant Breeding and Seed Science. 2014;67(1): 115-23.
53. Oliveira J, Fernandes V, Miranda C, Santos-Buelga C, Silva A, de Freitas V, Mateus N. Color properties of four cyanidin– pyruvic acid adducts. Journal of agricultural and food chemistry 2006; 54(18): 6894-903.
54. Alasalvar C, Al-Farsi M, Quantick PC, Shahidi F, Wiktorowicz R. Effect of chill storage and modified atmosphere packaging (MAP) on antioxidant activity, anthocyanins, carotenoids, phenolics and sensory quality of ready-to-eat shredded orange and purple carrots. Food Chemistry 2005 ;89(1): 69-76.
55. Bochi VC. Chemical characterization and biological activity of ceylon gooseberry (*Ddovyalis hebecarpa*) in different stages of maturation. Food Chemistry 2004; 164: 1-556.
56. Pomilio AB, Sproviero JF, Fernandez ME. Anthocyanins from argentine *Erythrina* species. Asoc Quim Argent An. 1971; 59(1): 29-33
57. de Rosso VV, Hillebrand S, Montilla EC, Bobbio FO, Winterhalter P, Mercadante AZ. Determination of anthocyanins from acerola (*Malpighia emarginata* DC.) and açai (*Euterpe oleracea* Mart.) by HPLC–PDA–MS/MS. Journal of Food Composition and Analysis. 2008; 21(4): 291-9.
58. Okamura N, Haraguchi H, Hashimoto K, Yagi A. Flavonoids in *Rosmarinus officinalis* leaves. Phytochemistry 1994;37 (5): 1463-6.
59. Carbone F, Preuss A, De Vos RC, D'AMICO EL, Perrotta G, Bovy AG, Martens S, Rosati C. Developmental, genetic and environmental factors affect the expression of flavonoid genes, enzymes and metabolites in strawberry fruits. Plant, Cell & Environment 2009;32(8): 1117-31.
60. Halbwirth H, Puhl I, Haas U, Jezik K, Treutter D, Stich K. Two-phase flavonoid formation in developing strawberry (*Fragaria*  $\times$  *ananassa*) fruit. Journal of Agricultural and Food chemistry 2006;54(4):1479-85.
61. Nayak CA, Srinivas P, Rastogi NK. Characterisation of anthocyanins from *Garcinia indica* Choisy. Food chemistry. 2010; 118(3): 719-24.
62. Cho KM, Ha TJ, Lee YB, Seo WD, Kim JY, Ryu HW, Jeong SH, Kang YM, Lee JH. Soluble phenolics and antioxidant properties of soybean (*Glycine max* L.) cultivars with varying seed coat colours. Journal of Functional Foods 2013;5(3): 1065-76.
63. Shimizu Y, Maeda K, Kato M, Shimomura K. Isolation of anthocyanin-related MYB gene, GbMYB2, from *Gynura bicolor* leaves. Plant biotechnology 2010; 27(5): 481-7.
64. Jordheim M, Skaar I, Lunder H, Andersen ØM. Anthocyanins from *Fuchsia* flowers. Natural product communications. 2011; 6(1): 35-40.
65. Du CT, Francis FJ. Anthocyanins of roselle (*Hibiscus sabdariffa*, L.). Journal of Food Science. 1973; 38(5): 810-2.
66. Woolfe JA. Sweet potato: an untapped food resource. Cambridge University Press; 1992.
67. Chen YB, Wu KS, Gu Y, Chen JZ. Research progress in the chemical constituents and pharmacological effects of lychee seeds. Chinese Journal of Infection 2007;14: 97-8.
68. Kowalczyk E, Krześniński P, Kura M, Szmigiel B, Blaszczyk J. Anthocyanins in medicine. Polish journal of pharmacology 2003; 55(5):699-702.
69. Yuan Y, Yang J, Yu X, Huang L, Lin S. Anthocyanins from buds of *Lonicera japonica* Thunb. var. *chinensis* (Wats.) Bak. Food research international 2014; 62: 812-8.
70. Saito N, Tatsuzawa F, Nishiyama A, Yokoi M, Shigihara A, Honda T. Acylated cyanidin 3-sambubioside-5-glucosides in *Matthiola incana*. Phytochemistry 1995; 38(4): 1027-32.
71. Liu LK, Chou FP, Chen YC, Chyau CC, Ho HH, Wang CJ. Effects of mulberry (*Morus alba* L.) extracts on lipid

- homeostasis in vitro and in vivo. *Journal of agricultural and food chemistry* 2009; 57(16): 7605-11.
72. Peng CH, Liu LK, Chuang CM, Chyau CC, Huang CN, Wang CJ. Mulberry water extracts possess an anti-obesity effect and ability to inhibit hepatic lipogenesis and promote lipolysis. *Journal of agricultural and food chemistry* 2011; 59(6): 2663-71.
  73. Simmonds NW. *The Evolution of the Bananas. The Evolution of the Bananas.* 1<sup>st</sup>ed.1962. [London] Longmans publishers; 1962.
  74. Reynertson KA, Wallace AM, Adachi S, Gil RR, Yang H, Basile MJ et al. Bioactive Depsides and Anthocyanins from Jaboticaba (*Myrciaria c auliflora*). *Journal of Natural Products* 2006; 69(8): 1228-30.
  75. Bao J, Cai Y, Sun M, Wang G, Corke H. Anthocyanins, flavonols, and free radical scavenging activity of Chinese bayberry (*Myrica rubra*) extracts and their color properties and stability. *Journal of Agricultural and Food Chemistry* 2005;53(6): 2327-32.
  76. Martin T, Villaescusa L, Sotto MD, Lucia A, Diaz AM. Determination of anthocyanic pigments in *Myrtus communis* berries. *Fitoterapia - Journal.* 1990; 61(1).
  77. Wu X, Prior RL. Systematic identification and characterization of anthocyanins by HPLC-ESI-MS/MS in common foods in the United States: fruits and berries. *Journal of agricultural and food chemistry* 2005; 53(7):2589-99.
  78. Phippen WB, Simon JE. Anthocyanins in basil (*Ocimum basilicum* L.). *Journal of Agricultural and Food Chemistry.* 1998; 46(5): 1734-8.
  79. Strack D, Busch E, Klein E. Anthocyanin patterns in European orchids and their taxonomic and phylogenetic relevance. *Phytochemistry* 1989; 28(8): 2127-39.
  80. Kim MK, Kim HA, Koh K, Kim HS, Lee YS, Kim YH. Identification and quantification of anthocyanin pigments in colored rice. *Nutrition Research and Practice.* 2008; 2(1): 46-9.
  81. Hosoki T, Hamada M, Kando T, Moriwaki R, Inaba K. Comparative study of anthocyanins in tree peony flowers. *Journal of the Japanese Society for Horticultural Science* 1991; 60(2): 395-403.
  82. Wang LS, Hashimoto F, Shiraishi A, Aoki N, Li JJ, Sakata Y. Chemical taxonomy of the Xibei tree peony from China by floral pigmentation. *Journal of plant research* 2004;117(1): 47-55.
  83. Choung MG, Choi BR, An YN, Chu YH, Cho YS. Anthocyanin profile of Korean cultivated kidney bean (*Phaseolus vulgaris* L.). *Journal of agricultural and food chemistry.* 2003; 51(24): 7040-3.
  84. An C, Ichinose Y, Yamada T, Tanaka Y, Shiraishi T, Oku H. Organization of the genes encoding chalcone synthase in *Pisum sativum*. *Plant molecular biology.* 1993; 21(5): 789-803.
  85. Merina AJ, Sivanesan D, Begum VH, Sulochana N. Antioxidant and hypolipidemic effect of *Plumeria rubra* L. in alloxan induced hyperglycemic rats. *Journal of Chemistry.* 2010; 7(1):1-5.
  86. Kelley DS, Rasooly R, Jacob RA, Kader AA, Mackey BE. Consumption of Bing sweet cherries lowers circulating concentrations of inflammation markers in healthy men and women. *The Journal of nutrition* 2006; 136(4):981-6.
  87. Garofulić IE, Dragović-Uzelac V, Jambrak AR, Jukić M. The effect of microwave assisted extraction on the isolation of anthocyanins and phenolic acids from sour cherry Marasca (*Prunus cerasus* var. Marasca). *Journal of Food Engineering* 2013; 117(4): 437-42.
  88. Macheix JJ, Fleuriet A. *Fruit phenolics.* CRC press; 1990 Mar 20.
  89. Vinson JA, Su X, Zubik L, Bose P. Phenol antioxidant quantity and quality in foods: fruits. *Journal of Agricultural and Food Chemistry* 2001; 49(11): 5315-21.
  90. Gil MI, García-Viguera C, Artés F, Tomás-Barberán FA. Changes in pomegranate juice pigmentation during ripening. *Journal of the Science of Food and Agriculture.* 1995; 68(1):77-81.
  91. Aviram M, Dornfeld L, Kaplan M, Coleman R, Gaitini D, Nitecki S, Hofman A, Rosenblat M, Volkova N, Presser D, Attias J. Pomegranate juice flavonoids inhibit low-density lipoprotein oxidation and cardiovascular diseases: studies in atherosclerotic mice and in humans. *Drugs under experimental and clinical research* 2002; 28(2-3): 49-62.
  92. Mazza G, Miniati E. *Anthocyanins in fruits, vegetables, and grains.* 1<sup>st</sup> edition: Boca Raton: CRC Press publishers; 1993.
  93. Santhakumar AB, Kundur AR, Sabapathy S, Stanley R, Singh I. The potential of anthocyanin-rich Queen Garnet plum juice supplementation in alleviating thrombotic risk under induced oxidative stress conditions. *Journal of Functional Foods* 2015; 14: 747-57.
  94. Lin LZ, Sun J, Chen P, Harnly JA. LC-PDA-ESI/MS n identification of new anthocyanins in purple Bordeaux radish (*Raphanus sativus* L. variety). *Journal of agricultural and food chemistry* 2011;59(12): 6616-27.
  95. Liu GL, Guo HH, Sun YM. Optimization of the extraction of anthocyanins from the fruit skin of *Rhodomyrtus tomentosa* (Ait.) Hassk and identification of anthocyanins in the extract using high-performance liquid chromatography-electrospray ionization-mass spectrometry (HPLC-ESI-MS). *International journal of molecular sciences* 2012; 13(5): 6292-302.
  96. Frøyttlog C, Slimestad R, Andersen ØM. Combination of chromatographic techniques for the preparative isolation of anthocyanins—applied on blackcurrant (*Ribes nigrum*) fruits. *Journal of Chromatography A* 1998; 825(1): 89-95.
  97. Wenzig EM, Widowitz U, Kunert O, Chrubasik S, Bucar F, Knauder E, Bauer R. Phytochemical composition and in vitro pharmacological activity of two rose hip (*Rosa canina* L.) preparations. *Phytomedicine* 2008 ;15(10): 826-35.
  98. Ding M, Feng R, Wang SY, Bowman L, Lu Y, Qian Y et al. Cyanidin-3-glucoside, a natural product derived from blackberry, exhibits chemopreventive and chemotherapeutic activity. *Journal of Biological Chemistry* 2006; 281(25): 17359-68.
  99. Elisía I, Hu C, Popovich DG, Kitts DD. Antioxidant assessment of an anthocyanin-enriched blackberry extract. *Food chemistry- journal* 2007;101(3): 1052-8.
  100. Mes PJ, Boches P, Myers JR, Durst R. Characterization of tomatoes expressing anthocyanin in the fruit. *Journal of the American Society for Horticultural Science* 2008;133(2):262-9.
  101. Tanchev SS, Ruskov PJ, Timberlake CF. The anthocyanins of Bulgarian aubergine (*Solanum melongena*). *Phytochemistry* 1970; 9(7): 1681-2.
  102. Brown CR, Wrolstad R, Durst R, Yang CP, Clevidence B. Breeding studies in potatoes containing high concentrations of anthocyanins. *American Journal of potato research* 2003; 80(4): 241-9.
  103. Zhang LL, Lin YM. Antioxidant tannins from *Syzygium cumini* fruit. *African Journal of Biotechnology* 2009;8(10).
  104. Abdel-Aal ES, Hucl P. Composition and stability of anthocyanins in blue-grained wheat. *Journal of Agricultural and food Chemistry* 2003; 51(8): 2174-80.
  105. Matsunaga N, Imai S, Inokuchi Y, Shimazawa M, Yokota S, Araki Y, Hara H. Bilberry and its main constituents have neuroprotective effects against retinal neuronal damage in



- vitro and in vivo. *Molecular nutrition & food research* 2009; 53(7): 869-77.
106. Gao R, Cao B, Hu Y, Feng Z, Wang D, Hu W, Chen J, Jie Z, Qiu H, Xu K, Xu X. Human infection with a novel avian-origin influenza A (H7N9) virus. *New England Journal of Medicine* 2013;368(20):1888-97.
107. Matchett MD, MacKinnon SL, Sweeney MI, Gottschall-Pass KT, Hurta RA. Blueberry flavonoids inhibit matrix metalloproteinase activity in DU145 human prostate cancer cells. *Biochemistry and Cell Biology*. 2005; 83(5):637-43.
108. Milbury PE, Vita JA, Blumberg JB. Anthocyanins are bioavailable in humans following an acute dose of cranberry juice. *The Journal of nutrition* 2010; 140(6): 1099-104.
109. Roggero JP, Coen S, Ragonnet B. High performance liquid chromatography survey on changes in pigment content in ripening grapes of Syrah. An approach to anthocyanin metabolism. *American Journal of Enology and Viticulture* 1986; 37(1): 77-83.
110. Tsuda T, Kato Y, Osawa T. Mechanism for the peroxynitrite scavenging activity by anthocyanins. *FEBS letters* 2000; 484(3): 207-10.

**Cite this article as:**

Ashya Shaik et al. A review on Anthocyanins: A promising role on phytochemistry and pharmacology. *Int. Res. J. Pharm.* 2018;9(1):1-9 <http://dx.doi.org/10.7897/2230-8407.0911>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.