



Research Article

SYNTHESIS AND CHARACTERIZATION OF NOVEL RUTIN DERIVATIVES WITH POTENTIAL ANTIOXIDANT PROPERTIES

Mahitab H. El Bishbishy ¹ and Mohammed Abdalla Hussein ^{2*}

¹Pharmacognosy Department, Faculty of Pharmacy, MSA University, 12585, Giza, Egypt

²Biochemistry Department, Faculty of Applied Medical Sciences, October 6 University, 12585, Giza, Egypt

*Corresponding Author Email: prof.husseinma@o6u.edu.eg

Article Received on: 03/10/18 Approved for publication: 02/11/18

DOI: 10.7897/2230-8407.0911253

ABSTRACT

Rutin, a flavonol glycoside isolated from the fruit peels of *Malus domestica* family Rosaceae, is well known to possess antioxidant activity. This research was conducted in order to synthesize and characterize rutin derivatives and evaluate their antioxidant activities. Four derivatives were synthesized namely, Rutin-oxy-5, 7, 4' acetic acid (**2**), Rutin -oxy- 5,7, 4' methyl benzoate amide (**3**), 2'', 2''', 2''''-rutinoxymethyl-3-amino-1 -benzo[5,6-a] pyrimidine-4-one (**4**) and 5'', 5''', 5''''-tri-p-methoxybenzylideneamino-6-rutinoxy-benzo[5,6-a]-pyrimidin-4(5H)-one (**5**). Their structures were elucidated using different spectral data (Mass, IR and ¹H, ¹³C NMR). The antioxidant activities of rutin and its derivatives were evaluated by reducing power, Fe⁺² chelating, DPPH and ferric thiocyanate (FTC) assays. In addition, the results were compared with natural and synthetic antioxidants, such as α -tocopherol, butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA) and trolox. Rutin and its derivatives were exhibited a strong reducing power, chelating activity on Fe²⁺ and free radical-scavenging. Antioxidant activity of rutin and its derivatives increased with increased concentrations. Total antioxidant activity of rutin, its derivatives and both standards decreased in the order of rutin > compound (5) > trolox > compound (2) > BHA > compound (4) > compound (3). This study showed that Rutin and its derivatives exhibited antioxidant activity in all tests and could be considered as a source of natural and synthetic antioxidants.

Keywords: *Malus domestica*, Rutin, Reducing power assay - Fe⁺² chelating assay - DPPH assay - Ferric thiocyanate assay.

INTRODUCTION

Rutin (3,3',4',5,7-pentahydroxyflavone-3-rhamnoglucoside; Figure 1) is a flavonoid of the flavonol-type that is widespread in the plant kingdom¹. Rutin has a wide range of pharmacological properties (e.g., antioxidative activity) that have been exploited in human medicine and nutrition. Conventionally, it is used as an antimicrobial, antifungal, and anti-allergic agent. However, current research has shown its multi-spectrum pharmacological benefits for the treatment of various chronic diseases, such as cancer, diabetes, hypertension, and hypercholesterolemia². Rutin exerts anti-inflammatory effects in ultraviolet β -irradiated mouse skin by inhibiting the expression of cyclooxygenase-2 and inducible nitric oxide synthase³.

On our going medicinal chemistry research program we found that pyrimidine derivatives exhibit potent anticancer⁴⁻¹² and antibacterial¹³⁻¹⁵ activities. Also, pyrimidine derivatives were found to possess several pharmacological properties, including antiviral¹⁶, antifungal¹⁷, hypoglycemic¹⁸ and diuretic¹⁹ activities. As an extension of our studies on the synthesis of some new biologically active heterocyclic compounds^{18, 19}. In this manuscript, we are reporting the synthesis of combining form of these two structural features into one molecule might produce compounds with promising antioxidant effects.

MATERIALS AND METHODS

Plant Material

Apple fruits (*Malus domestica* Borkh. nom. illeg., cultivar Anna, family Rosaceae) were collected from the Faculty of Agriculture

eco-station, Faculty of Agriculture Cairo University, and authenticated by the Vegetables and fruits department, Faculty of Agriculture, Cairo University. Herbarium voucher specimen number are kept at Faculty of Pharmacy, MSA University number (2017-MSA-11). Their peels were cleaned, scraped off, then dried in shade, in a clean dust free environment at room temperature for several days and then grinded as powder. The powder obtained was stored in an air-tight container.

Chemicals and Reagents

All chemicals were purchased from Sigma Chemicals Co., St. Louis, MO, USA. Precoated TLC plates silica gel 60 GF₂₅₄ (20 x 20 cm) (E-Merck), polyamide for column chromatography 6, (Fluka) and sephadex LH-20 for column chromatography (Pharmazia)

Apparatus

Rotatory evaporator, Heidolph, (G. Switzerland) was used for distilling off the different solvents. Melting points were determined in open capillaries and are uncorrected. ¹H- and ¹³CNMR spectra were recorded in DMSO-d₆ and CD₃OD, respectively, on a JEOL ECA-500 (¹H, 500 MHz and ¹³C, 125 MHz), National Research Center, Giza, Egypt. UV-visible spectrophotometer, Shimadzu UV (P/N 204-58000) was used for all spectrophotometric assays. Melting points were determined on Gallenkamp melting point apparatus and are uncorrected. The infrared (IR) spectra were recorded on Shimadzu MR 470 infrared spectrophotometer using the KBr pellets.

Extraction and isolation

Three hundred and twenty grams of the powdered apple peels were macerated in 70% aqueous ethanol till exhaustion. The hydro-ethanolic extracts were concentrated under reduced pressure at a temperature not exceeding 50°C to yield 47 g dark brown dried extract. The dried residue was suspended in water and partitioned successively between water and petroleum ether (5×50ml), dichloromethane (5×50ml), ethyl acetate (5×50ml) and n-butanol (5×50ml). The solvents were distilled off to yield respective fractions of 8.3 g, 6.9 g, 12.4 g and 5.2 g. The ethyl acetate fraction was further fractionated over a column chromatography packed with 200 gm polyamide as a stationary phase. The elution process started with distilled water, followed by gradient decrease in the polarity by adding methanol, starting with 10% up to 100% methanol. Fractions were collected each 25 ml and monitored by TLC on silica gel pre-coated plates using (ethyl acetate-methanol-water 100:16.5:13.5 v/v/v) as a mobile phase. The spots were visualized under UV light before and after exposure to ammonia vapors and spraying with aluminum chloride reagent. Similar fractions were pooled and separately concentrated under reduced pressure at a temperature not exceeding 50°C. The water-methanol 7:3 eluted sub-fraction was subjected to sephadex column chromatography and further eluted with a stepwise gradient elution of water and methanol which resulted in the isolation of Compound 1 (31 mg). Its structure was elucidated by spectroscopic means.

Synthesis of rutin derivatives

Synthesis of Rutin-oxy-5, 7, 4' acetic acid (2)

A mixture of rutin (0.01 mol) and 0.03M NaOH in ethanol/water (2:8 v/v) was refluxed for 4hr. After cooling 0.03 M chloroacetic acid was added drop-wise, the solid obtained was recrystallized from ethanol to give (2).

Synthesis of Rutin -oxy- 5,7, 4' methyl benzoate amide (3)

To a solution of Rutin-oxy-5, 7, 4' acetic acid (2) (0.01 mol) and methyl anthranilate (6.93 g, 0.03 mol) in xylene (50 ml), phosphorus trichloride (3 ml) was added. The reaction mixture was heated under reflux for 3–4 h. The crude product was recrystallized from ethanol to give (3).

Synthesis of 2'', 2''', 2''''-rutinoxymethyl-3-amino-1 -benzo[5,6-a]pyrimidine-4-one (4)

A mixture of Rutin -oxy- 5, 7, 4' methyl benzoate amide (3) (0.01 mol) and hydrazine hydrate (95 %) (0.15 mol) were dissolved in n-butanol (30 ml) and refluxed for 3–5 h. The solvent was concentrated and the residue was recrystallized from ethanol to give (4).

Synthesis of 5'', 5''', 5''''-tri-p-methoxybenzylideneamino-6-rutinoxy-benzo [5, 6-a]-pyrimidin-4(5H)-one (5)

A mixture of 2'', 2''', 2''''-rutinoxymethyl-3-amino-1 -benzo[5,6-a]pyrimidine-4-one (4) (0.01 mol) and p-anisaldehyde (0.03 mol) in n-butanol was refluxed for 4hr. The reaction mixture was poured onto ice water and the solid obtained was recrystallized from dioxane to give (5).

Biological Investigations

Estimation of antioxidant activity

The reducing power of compounds 1-5 were measured according to the method of Oyaizu²⁰. α -tocopherol, BHA and BHT were

used as standard antioxidants. Higher absorbance of the reaction mixture indicated greater reducing power.

Determination of Fe⁺² chelating activity

The chelating activity of compounds (1-5) on ferrous ions (Fe⁺²) were measured according to the method of Decker and Welch²¹. The chelating activity of compounds (1-5) on Fe⁺² were compared with that of EDTA at a level of 0.037 mg/ml. Chelating activity was calculated using the following formula:

$$\text{Chelating activity (\%)} = [1 - (\text{Absorbance of sample} / \text{Absorbance of control}) \times 100]$$

Control test was performed without addition of compounds 1-5.

Determination of DPPH assay

The free radical-scavenging activity of compounds 1-5 were measured with 1, 1-diphenyl-2-picrylhydrazil (DPPH[•]) using the slightly modified methods of Brand-Williams *et al.*²² and Takashira and Ohtake²³. The percent inhibition activity was calculated using the following equation:

$$\text{Inhibition activity (\%)} = [(A_0 - A_1) / A_0 \times 100]$$

Where A₀ is the absorbance of the control reaction and A₁ is the absorbance in the presence of compounds 1-5 samples

Determination of FTC

The antioxidant activity was determined according to the thiocyanate method of Osawa and Namiki²⁴ with slight modifications. The solutions without added compounds 1-5 or standards were used as the control. The inhibition of lipid peroxidation in percentage was calculated by the following equation.

$$\text{Inhibition activity (\%)} = [(A_0 - A_1) / A_0 \times 100]$$

Where A₀ is the absorbance of the control reaction and A₁ is the absorbance in the presence of compounds 1-5 or standards.

Statistical analysis

All the grouped data were statistically evaluated with SPSS/15 software²⁵. Hypothesis testing methods included one-way analysis of variance (ANOVA). P < 0.05 was considered to indicate statistical significance. All the results were expressed as mean \pm SD for six replicates.

RESULTES AND DISCUSSION

Structure elucidation of the isolated rutin (1)

Yellowish powder obtained from ethyl acetate fraction. The structure of isolated rutin with melting point of 230-232°C is presented in Figure 1. IR spectrum of rutin showed that 3365, 3210 (O-H stretching), 1510(C=O groups) and 1390(C-OH vibrations); ¹HNMR spectrum of rutin displayed the characteristic 3.02-3.6 (m, 12H of sugar moieties), 4.3 (d, J=1.15Hz, 1H-Rham), 0.95 (3H, d, J=6Hz, CH₃-Rham), 6.1 (4H, d, J=7.8Hz, H-1 Glu), 5.3 (1H, d, J=2Hz H-6), 6.2 (1H, d, J=2Hz, H-8), 6.7 (1H, d, J=8Hz, H-5'), 6.99, 7.4, 6.93 (1H, , H-2', H-6'). The ¹³CNMR spectrum showed the presence of specific signals (aromatic carbon, aliphatic chain and methoxy group) in agreement with the proposed structure of rutin. Also, the ¹³CNMR spectrum showed the presence of 161.73 (C-2), 133.8 (C-3), 177.9(C-4), 164.6 (C-5), 104.49(C-6), 168.5(C-7), 94.12 (C-8), 156.9(C-9), 101.7(C-10), 121.7(C-1'), 116.3(C-2'), 145.6(C-3'), 148.6(C-4'), 122.1(C-5'), 126.2(C-6'), 99.2(C-1''), 68.8(C-2''), 76.4(C-3''), 74.6(C-4''), 72.3(C-5''), 71.06(C-6''), 40.1(C-1'''),

40.00(C-2'''), 39.8(C-3'''), 35.18(C-4'''), 32.8(C-5'''), 24.7 (C-6''').

Structure elucidation of Rutin derivatives compounds (2-5)

Compound 2

Preparation of the target compound *Rutin-oxy-5, 7, 4' acetic acid* (2) were achieved by the route depicted in Scheme 1. The structure of compound (2) was proved on the basis of elemental analyses, IR, ¹H NMR and ¹³C NMR spectral data. The IR spectrum of compound (2) showed bands at 3400 (O-H stretching), 2916, 2819 (CH₂-stretching), 2664 (C-H bonding) and 1701, 1678, 1585, 1541(4C=O groups). ¹H NMR spectrum of compound (2) in (DMSO_d₆) exhibited signals at 3.2-3.5(m,12H of sugar moieties), 3.70(d, J=1.15Hz, 1H-Rham), 1.52(3H,d,J=6Hz, CH₃-Rham), 6.3(4H,d, J=7.8Hz, H-1 Glu), 6.31 (1H,d,=2Hz H-6), 7.1 (1H, d, J=2Hz, H-8), 7.4 (1H, d, J=8Hz, H- 5'), 7.45, 7.46, 7.5 (1H, , H-2', H-6'), 10.3 (s,1H,carboxylic proton). ¹³C NMR spectrum of compound (2) in (DMSO_d₆) exhibited signals at 156.20 (C-2), 124.84(C-1'), 114.6(C-2''), 140.42(C-3''), 145.5(C-4'), 127.82(C-5''), 129.79(C-6'), 101.0(C-1'''), 48.6(C-2''), 40.7(C-3''), 40.5(C-4''), 40.71(C-5''), 40.34(C-6''), 40.2(C-1'''), 39.9(C-2'''), 39.5(C-3'''), 35.18(C-4'''), 32.8(C-5'''), 24.7 (C-6''').

Compound 3

The IR spectrum of compound 3 showed bands at 3390 (O-H stretching) and 1725, 1701, 1690, 1662, 1624, 1600, 1535(7C=O groups). ¹H NMR spectrum of compound 3 in (DMSO_d₆) exhibited signals at 3.12-3.6, 3.80, 1.04 and 2.46 indicated that the presence of sugar moieties, 1H-Rham, CH₃-Rham, H-1 Glu. and COCH₃, respectively, as well as the absence of carboxylic protons. ¹³C NMR spectrum of compound 3 in (DMSO_d₆) exhibited signals at 182.47 (C-2), 123.37(C-1'), 119.47(C-2''), 147.26(C-3''), 140.5(C-4'), 125.4(C-5''), 130.0(C-6'), 95.30(C-1'''), 70.07, 68.86, 56.9(3 methoxy groups), 40.12(C-2''), 39.95(C-3''), 39.7(C-4''), 38.7(C-5''), 35.47(C-6''), 33.65(C-1'''), 34.60(C-2'''), 30.54(C-3'''), 29.8(C-4''), 32.8(C-5''), 18.35 (C-6''').

Compound 4

The IR spectrum of compound 4 showed bands at 3398 (O-H stretching), 3178, 3274, 3124 cm⁻¹ (NH₂) and 1732, 1658, 1690, 1620 (4C=O groups). ¹H NMR spectrum of compound 4 in (DMSO_d₆) exhibited signals at 3.3-3.7, 3.97, 1.13, 6.73 and 6.73, 7.00 indicated that the presence of sugar moieties, 1H-Rham, CH₃-Rham, H-1 Glu. and NH₂ respectively, as well as the absence of COCH₃ group. ¹³C NMR spectrum of compound 4 in (DMSO_d₆) exhibited signals at 200.9 (C-2), 112.6(C-1'), 108.24(C-2''), 135.7(C-3''), 134.9(C-4'), 125.6(C-5''), 116.47(C-6'), 61.3(C-1'''), 51.8(C-2''), 56.3(C-3''), 39.69(C-4''), 29.7(C-5''), 23.00(C-6''), 21.3(C-1'''), 24.52(C-2'''), 20.44(C-3'''), 19.7(C-4''), 17.7(C-5''), 14.01 (C-6''').

Compound 5

The IR spectrum of compound 5 showed bands at 3390 (O-H stretching), 2923, 2870 (CH₂-stretching), 2568 (C-H bonding), 1700, 1635, 1596, 11562(4C=O groups) and 1350 (C-OH vibrations) and indicated that the absence of NH₂ group. ¹H NMR spectrum of compound 5 in (DMSO_d₆) exhibited signals at 3.4-3.7, 4.1, 1.2 and 6.98 indicated that the presence of sugar moieties, 1H-Rham, CH₃-Rham. and H-1 Glu. as well as the absence of NH₂ signal, respectively. Also, ¹H NMR spectrum indicated that the absence of signal at 6.2 [s, 1H, N=CH, proton]. ¹³C NMR spectrum of compound 5 in (DMSO_d₆) exhibited signals at 176.03 (C-2), 120.9(C-1'), 96.6(C-2''), 143.7(C-3''),

130.5(C-4'), 116.9(C-5'), 115.8(C-6'), 71.9(C-1'''), 56.17(C-2''), 53.9(C-3''), 40.15(C-4''), 39.98(C-5''), 39.82(C-6''), 27.18(C-1'''), 24.52(C-2'''), 23.5 (C-5''), 22.65 (C-5'') and 20.7(3C-5'') indicated that the presence of 3 double bond at [N=CH], 18.98 (C-6''').

Antioxidant activities

It is well known from the literature that the tested rutin ant it's derivatives exhibit a wide range of biological activities. So, it was of interest to design new compounds containing both these biologically active phenolic moieties and to study their antioxidant activities.

Figure 1 shows the reducing power of compounds 1-5. The reducing power of compounds 1-5 increased with increasing concentration. Based on a comparison of the absorbance at 700 nm, the reducing power of compounds 1 and 5 at a concentration of 40 µg/ml were nearly similar to that of BHT. This indicates that compounds 1 and 5 were electron donors and could also react with free radicals, converting them to more stable products and terminate the radical chain reaction. Also, 120 µg/ml compounds 1-5 are the best concentration which exhibits the most reducing power. In the reducing power assay, the presence of reductants (antioxidants) resulted in the reduction of the Fe⁺³/ferricyanide complex to the ferrous form (Fe⁺²) table 1. The amount of Fe⁺² complex can therefore be monitored by measuring the formation of Perl's Prussian Blue at 700 nm. The reducing power of compounds 1-5 and both standards decreased in the order of compound 1 > compound 5 > BHT > compound 2 > compound 4 > α-tocopherol > BHA > compound 3. Table 2 shows the chelating effect of compounds 1-5. All samples at 1.50 mg/ml concentration showed the best chelating effect 48.87, 33.5, 20.8, 34.3 and 43.89% on ferrous ions at an incubation time of 60 min, respectively. The chelating activity of samples increased with increasing incubation times with FeCl₂. However, the chelating activity of compounds 2 of 1.25 mg/ml (30 %) was lower than EDTA at 0.037 mg/ml (43.67 %) for an incubation time of 90 min. This indicates that the chelation property of the compounds 1-5 on Fe⁺² ions may afford protection against oxidative damage. The results in table 2, indicated that a significant property of compounds 1-5 is its capability for blocking the oxidative activity of systems with transition metal ion (Fe⁺²/Fe⁺³) that play an essential role in the formation of reactive oxygen species in Fenton's reactions.

The DPPH* radical-scavenging effects of compounds 1-5 are presented in Fig. 2 and showed appreciable free radical-scavenging activities. The free radical-scavenging activity of compounds 1-5 were compared to trolox, as a synthetic antioxidant. Compounds 1-5 of 6 mg/ml had the highest radical-scavenging activity when compared with 0.06 mg/ml trolox. The effects of 100 mg/l of compounds 1-5 on peroxidation of linoleic acid emulsion are shown in Fig. 2.

Compound 1(0.06mg/ml) showed higher antioxidant activity when compared to trolox, BHA, and BHT. Total antioxidant activity of compounds 1-5 and both standards decreased in the order of compound 1 > compound 5 > trolox > compound 4 > compound 2 > compound 3. Free radicals are known to be a major factor in biological damages, and DPPH* has been used to evaluate the free radical-scavenging activity of natural antioxidants²⁶. DPPH*, which is a molecule containing a stable free radical with a purple color, changes into a stable compound with a yellow color by reacting with an antioxidant which can donate an electron to DPPH*. In such case, the purple color typical of the free DPPH* radical decays, a change which can be followed either spectrophotometrically (517 nm). The proton radical-scavenging action is known as an important mechanism of antioxidation. 1, 1-Diphenyl-2-picrylhydrazil (DPPH*) is used as

a free radical to evaluate the antioxidative activity of some natural sources²⁷. The DPPH[•] radical-scavenging effects of compounds 1-5 are presented in Fig. 3. From these results, it can be stated that compounds 1 and 5 have the ability to scavenge free radicals and could serve as a strong free radical inhibitor or scavenger according to trolox. Many attempts at explaining the structure–activity relationships of some phenolic compounds have been reported in the literature. It has been reported that the antioxidant activity of phenolic compounds may result from the neutralization of free radicals initiating oxidation processes, or from the termination of radical chain reactions, due to their

hydrogen donating ability²⁸. It is also known that the antioxidant activity of phenolic compounds is closely associated with their structures, such as substitutions on the aromatic ring and side chain structure. Their accessibility to the radical center of DPPH[•] could also influence the order of the antioxidant power. Free radical scavenging activity of phenolic compounds is believed to be influenced by the number and position of phenolic hydrogen in their molecules²⁹. It is also proposed that the higher antioxidant activity of compounds 1-5 is related to the presence of hydroxyl groups³⁰.

Table 1: Reducing power of compounds 1-5, BHA, BHT and α -tocopherol

		Concentrations				
Compounds		40 ug/ml	60 ug/ml	80 ug/ml	100 ug/ml	120 ug/ml
Absorbance (700 nm)	1	0.43±0.02 ^{Aa}	0.45±0.02 ^{Aa}	0.54±0.01 ^A	0.62±0.02 ^A	0.76±0.02 ^A
	2	0.28±0.01 ^{Ba}	0.32±0.02 ^B	0.42±0.04 ^{Ba}	0.50±0.03 ^B	0.51±0.03 ^B
	3	0.19±0.03 ^{Ca}	0.21±0.01 ^{Ca}	0.32±0.02 ^{Cb}	0.36±0.02 ^{Cb}	0.40±0.02 ^C
	4	0.26±0.02 ^B	0.35±0.02 ^{Ba}	0.37±0.01 ^{Ca}	0.44±0.04 ^{Bb}	0.48±0.04 ^{Bb}
	5	0.46±0.01 ^A	0.53±0.03 ^{Aa}	0.58±0.03 ^{Aa}	0.64±0.03 ^{Ab}	0.69±0.02 ^{Ab}
	α -tocopherol	0.33±0.05 ^{Ba}	0.34±0.01 ^{Ba}	0.39±0.02 ^C	0.46±0.02 ^{Bb}	0.45±0.01 ^{Cb}
	BHT	0.45±0.03 ^{Aa}	0.48±0.02 ^{Aa}	0.46±0.04 ^{Ba}	0.50±0.01 ^{Bb}	0.53±0.04 ^{Bb}
	BHA	0.25±0.04 ^{Ba}	0.27±0.03 ^{Ca}	0.34±0.02 ^{Cb}	0.39±0.01 ^{Cb}	0.41±0.02 ^C

Values are given as mean \pm SD for six replicates


Data followed by the same letter are not significantly different at $P \leq 0.05$. Small letters are used for comparison between the means within the columns. Capital letters are used to compare means within the rows.

Table 2: Chelating effects of different concentrations of compounds 1-5 on Fe²⁺ ions at different incubation times with FeCl₂

		Time (min)				
Compounds		10 min.	30 min.	60 min.	90 min.	120 min.
Chelating activity (%)	1	15.50 \pm 1.5 ^A	30.98 \pm 1.5 ^A	48.87 \pm 3.18 ^A	53.30 \pm 2.97 ^{Aa}	53.3 \pm 4.3 ^{Aa}
	2	10.38 \pm 0.98 ^B	25.48 \pm 2.45 ^B	33.50 \pm 2.04 ^{Ba}	36.09 \pm 2.77 ^a	36.99 \pm 3.25 ^a
	3	6.72 \pm 0.67	12.76 \pm 2.14	20.85 \pm 3.27 ^a	22.11 \pm 3.00 ^a	23.65 \pm 2.61 ^a
	4	9.86 \pm 0.9 ^B	21.50 \pm 2.71 ^B	34.30 \pm 3.08 ^B	44.28 \pm 4.22 ^a	46.90 \pm 2.59 ^a
	5	18.11 \pm 2.06 ^A	29.96 \pm 2.76 ^A	43.89 \pm 4.07 ^A	49.11 \pm 3.80 ^{Aa}	50.25 \pm 3.97 ^{Aa}

Values are given as mean \pm SD for six replicates

Data followed by the same letter are not significantly different at $P \leq 0.05$. Small letters are used for comparison between the means within the columns. Capital letters are used to compare means within the rows.



Scheme 1: Rationalized formation of compounds 1-5




Figure 1: Chemical structure of rutin




Fig. 2: Scavenging activities of different concentrations of compounds 1-5 and Trolox against the 1, 1-diphenyl-2-picryl-hydrazil (DPPH) radical




Fig. 3: Total antioxidant activities of compounds 1-5, trolox and BHA, (100 mg/l concentration) on peroxidation of linoleic acid emulsion

CONCLUSION

In conclusion, rutin when combining with quinazoline derivatives exhibited promising antioxidant activity. More studies are needed to prove their medicinal and biological importance which may pave the way for possible therapeutic applications.

REFERENCES

- Hosseinzadeh, H., Nassiri-Asl, M. Review of the protective effects of rutin on the metabolic function as an important

- Sharma, S., Ali, A., Ali, J., Sahni, J.K., Baboota, S. Rutin: therapeutic potential and recent advances in drug delivery. *Expert Opin Investig Drugs.* 2013; 22: 1063-79. <https://doi.org/10.1517/13543784.2013.805744>
- Choi, K.S., Kundu, J.K., Chun, K.S., Na, H.K., Surh, Y.J. Rutin inhibits UVB radiation-induced expression of COX-2 and iNOS in hairless mouse skin: p38 MAP kinase and JNK as potential targets. *Arch Biochem Biophys.* 2014; 559: 38-45. <https://doi.org/10.1016/j.abb.2014.05.016>

4. Ismail, M.M.F., Ghorab, M.M., Noaman, E., Ammar, Y.A., Heiba, H.I., Sayed, M.Y. Novel synthesis of pyrrolo[2,3-d]pyrimidines bearing sulfonamide moieties as potential antitumor and radioprotective agents. *Arzneimittel-Forschung (Drug Research)* 2006; 56(4):301–308.
5. Ghorab, M.M., Noaman, E., Ismail, M.M.F., Heiba, H.I., Ammar, Y.A., Sayed, M.Y. Novel antitumor and radioprotective sulfonamides containing pyrrolo[2,3-d]pyrimidines. *Arzneimittel-Forschung (Drug Research)*. 2006; 56(6):405–413.
6. Ghorab, M.M., Ragab, F.A., Noaman, E., Heiba, H.I., Galal, M. Synthesis of certain new thieno[2,3-d]pyrimidines as potential antitumor and radioprotective agents. *Arzneimittel-Forschung (Drug Research)*. 2006; 56(7):553–560.
7. Heiba, H.I., Ragab, F.A., Noaman, E., Ghorab, M.M., Galal, M. Synthesis of some novel sulphur containing triazolothienopyrimidines and biscompounds as possible antitumor and radioprotective agents. *Arzneimittel-Forschung (Drug Research)*. 2006; 56(8):593–599.
8. Ghorab, M.M., Osman, A.N., Noaman, E., Heiba, H.I., Zaher, N.H. The synthesis of some new sulfur heterocyclic compounds as potential radioprotective and anticancer agents. *Phosphorus Sulphur Silicon*. 2006;181:1935–1950. <https://doi.org/10.1080/10426500500544014>
9. Ghorab, M.M., Osman, A.N., Noaman, E., Heiba, H.I., Zaher, N.H. The Utility of isothiocyanato thiophenes in synthesis of thieno[2,3-d]pyrimidine derivatives as possible radioprotective and anticancer agents. *Phosphorus Sulphur Silicon*. 2006; 181: 1983–1996. <https://doi.org/10.1080/10426500500544238>
10. Chauhan, P.M.S, Martins, C.J.A, Horwell, D.C. Syntheses of novel heterocycles as anticancer agents. *Bioorg Med Chem*. 2005; 13:3513–3518. <https://doi.org/10.1016/j.bmc.2005.02.055>
11. Cocco, M.T., Congiu, C., Lilliu, V., Onnis, V. Synthesis and in vitro antitumoral activity new hydrazinopyrimidine-5-carbonitrile derivatives. *Bioorg Med Chem*. 2006; 14:366–372. <https://doi.org/10.1016/j.bmc.2005.08.012>
12. Amr, A.E., Mohamed, A.M., Mohamed, S.F., Abdel-Hafez, N.A., Hammam, A.G. Anticancer activities of some newly synthesized pyridine, pyrane, and pyrimidine derivatives. *Bioorg Med Chem*. 2006; 14:5481–5488. <https://doi.org/10.1016/j.bmc.2006.04.045>
13. Abdel-Gawad, S.M., El-Gaby, M.S.A., Heiba, H.I., Ali, H.M., Ghorab, M.M. Synthesis and radiation stability of some new biologically active hydroquinoline and pyrimido[4,5-b]quinoline derivatives. *J Chin Chem Soc*. 2005;52:1227–1236. <https://doi.org/10.1002/jccs.200500177>
14. El-Gaby, M.S., Abdel Gawad, S.M., Ghorab, M.M., Heiba, H.I., Ali, H.M. Synthesis and biological activity of some novel thieno[2,3-d]quinoline, quinolino[3,2:4,5]thieno[3,2-d]pyrimidine and pyrido[2,3:4,5]thieno[2,3-d]quinoline derivatives. *Phosphorus Sulphur Silicon*. 2006;181: 279–297. <https://doi.org/10.1080/104265090970322>
15. Selvi, S.T., Nadarajm V., Mohan, S., Sasi, R., Hema M. Solvent free microwave synthesis and evaluation of antimicrobial activity of pyrimido[4,5-b]- and pyrazolo[3,4-b]quinoline. *Bioorg Med Chem*. 2006; 14: 3896–3903. <https://doi.org/10.1016/j.bmc.2006.01.048>
16. Gawin, R., De Clercq, E., Naesens, L., Stawinska, M.K. Synthesis and antiviral evaluation of acyclic azanucleosides developed from sulfanilamide as a lead structure. *Bioorg Med Chem*. 2008;16(18):8379–89. <https://doi.org/10.1016/j.bmc.2008.08.041>
17. Ezabadi, I.R., Camoutsis, C., Zoumpoulakis, P., Geronikaki, A., Sokovic, M., Glamocilija, J. Sulfonamide-1,2,4-triazole derivatives as antifungal and antibacterial agents: Synthesis, biological evaluation, lipophilicity, and conformational studies. *Bioorg Med Chem*. 2008; 16(3):1150–61. <https://doi.org/10.1016/j.bmc.2007.10.082>
18. Hussein MA. Synthesis of some novel triazoloquinazolines and triazinoquinazolines and their evaluation for anti-inflammatory activity. *Medicinal Chemistry Research*. 2012; 21(8):1876–1886 <https://doi.org/10.1007/s00044-011-9707-0>
19. Hussein MA. Synthesis and Biochemical Evaluation of Some Novel Anti-inflammatory Quinazolines. *International Journal of Organic and Bioorganic Chemistry*. 2011; 1: 12-20.
20. Oyaizu M. Studies on products of browning reaction prepared from glucose amine. *Jpn J Nutr* 1986; 44:307–314. <https://doi.org/10.5264/eyogakuzashi.44.307>
21. Decker, E.A., Welch, B. Role of ferritin as a lipid oxidation catalyst in muscle food. *J Agr Food Chem* 1990; 38:674–683 <https://doi.org/10.1021/jf00093a019>
22. Brand-Williams, W., Cuvelier, M.E., Berset, C. Use of a free radical method to evaluate antioxidant activity. *Lebensm Wiss Technol* 1995; 28:25–32 [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)
23. Takashira, M., Ohtake, Y. A new antioxidative 1,3-benzodioxole from *Melissa officinalis*. *Planta Med* 1998; 64:555–563 <https://doi.org/10.1055/s-2006-957513>
24. Osawa, T., Namiki, N. A novel type of antioxidant isolated from leaf wax of *Eucalyptus* leaves. *Agric Biol Chem* 1981; 45:735–742 <https://doi.org/10.1080/00021369.1981.10864583>
25. SPSS. (SPSS 15, Inc., Chicago, IL, USA). 2012.
26. Yokozawa, T., Chen, C.P., Dong, E., Tanaka, T., Nonaka, G.I., Nishioka, I. Study on the inhibitory effect of tannins and flavonoids against the 1,1-diphenyl-2-picrylhydrazyl radical. *Biochem Pharmacol* 1998; 56:213–222. [https://doi.org/10.1016/S0006-2952\(98\)00128-2](https://doi.org/10.1016/S0006-2952(98)00128-2)
27. Chung, Y., Chen, S., Hsu, C., Chang, C., Chou, S. Studies on the antioxidative activity of *Graptopetalum paraguayense* E. Walther. *Food Chem* 2005; 91:419–423 <https://doi.org/10.1016/j.foodchem.2004.06.022>
28. Baumann, J., Wurn, G., Bruchlausen, F.V. Prostaglandin synthase inhibiting O₂-radical scavenging properties of some flavonoids and related phenolic compounds. *N-Ss Arch Pharmacol* 1979; 308:R27–R39.
29. Rice-Evans, C.A., Miller, N.J., Paganga, G. Structure antioxidant activity relationship of flavonoids and phenolic acids. *Free Radic Biol Med* 1996; 20:933–956 [https://doi.org/10.1016/0891-5849\(95\)02227-9](https://doi.org/10.1016/0891-5849(95)02227-9)
30. Cao, G., Sofic, E., Prior, R.L. Antioxidant and prooxidant behaviour of flavonoids: structure–activity relationships. *Free Rad Biol Med* 1997; 22(5):749–760 [https://doi.org/10.1016/S0891-5849\(96\)00351-6](https://doi.org/10.1016/S0891-5849(96)00351-6)

Cite this article as:

Mahitab H. El Bishbishy and Mohammed Abdalla Hussein. Synthesis and characterization of novel rutin derivatives with potential antioxidant properties. *Int. Res. J. Pharm.* 2018;9(11):31-36 <http://dx.doi.org/10.7897/2230-8407.0911253>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.