



## Research Article

### **IN-VITRO FREE RADICAL SCAVENGING ACTIVITY OF AERIAL PARTS OF ETHANOLIC EXTRACT OF *CORDIA OBLIQUA***

S. Sivakrishnan<sup>1\*</sup>, M. Swamivelmanickam<sup>2</sup>

<sup>1</sup>Assistant Professor, Department of Pharmacy, FEAT, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India

<sup>2</sup>Associate Professor, Department of Pharmacy, Annamalai University, Annamalai Nagar Tamil Nadu, India

\*Corresponding Author Email: sivaadm82@gmail.com

Article Received on: 20/06/19 Approved for publication: 28/07/19

**DOI: 10.7897/2230-8407.1008253**

#### ABSTRACT

Antioxidants are an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the body. Plant sources contain a variety of antioxidants. The ethanolic extract of *Cordia obliqua* was examined for DPPH ( $\alpha, \alpha$ -diphenyl- $\beta$ -picrylhydrazyl) radical scavenging activity, Superoxide anion, Nitric oxide and Hydroxyl radical scavenging activity with reference standard Rutin, Quercetin and Ascorbate respectively through *in vitro* models. Ethanolic extract of *Cordia obliqua* showed significant free radical scavenging activity than that of various standards. The radical scavenging activity was found to be concentration dependent manner. Ethanolic extract of *Cordia obliqua* showed strong scavenging activity against free radical compared to various standards. These *in vitro* assays indicate that this plant extract is a better source of natural antioxidant, which might be helpful in preventing the progress of various oxidative stresses.

**Keywords:** *Cordia obliqua*, Antioxidants, Rutin, Natural, Ascorbate, Ethanol

#### INTRODUCTION

Antioxidants are important in the prevention of human diseases. Oxygen free radicals induce damage due to peroxidation to bio membranes and also to DNA, which leads to tissue damage, thus cause occurrence of a number of diseases<sup>1</sup>. Antioxidants neutralize the effect of free radicals through different ways and may prevent the body from various diseases. A variety of free radical scavenging antioxidants is found in plant sources like aerial parts, fruits, vegetables, leaves, seeds, pods, etc<sup>2</sup>. *Cordia obliqua* Willd.<sup>3-7</sup> also known as Clammy cherry, is a flowering plant species in the genus *Cordia* belonging to the family Boraginaceae. It is found worldwide, mostly in warmer parts of India and Ceylon. Traditionally *Cordia obliqua* is used for cooling effects, anthelmintic, antimicrobial<sup>8</sup>, expectorant<sup>9,10</sup>, diuretic and it is used as treatment of diseases of spleen and leprosy. Our literature survey revealed that the antioxidant activity of various extracts from the whole plant of *Cordia obliqua* was not investigated; hence these activities have been investigated in the present study.

#### MATERIALS AND METHODS

##### Collection and Identification of *Cordia obliqua*

*Cordia obliqua* was collected from B. Maduvangarai, Chidambaram Taluk, Cuddalore District, Tamil Nadu, India.

Taxonomic identification was made from Botanical Survey of Medicinal Plants Unit Siddha, Government of India, Palayamkottai.

##### Extraction of *Cordia obliqua*

The leaves of *Cordia obliqua* were dried under shade, segregated, pulverized by a mechanical grinder and passed through a 40 mesh sieve. The powdered plant materials were stored in an air-tight container. The above powdered components were continuously extracted with ethanol in a Soxhlet apparatus using a 24 hours continuous hot percolation method. The extract was concentrated on a rotary evaporator and subjected to freeze drying in a lyophilizer until a dry powder was obtained<sup>11,12</sup>.

##### Evaluation of Antioxidant activity by *in vitro* Techniques

These are the following methods was used to evaluate the antioxidant activity by *in vitro* methods: The effect of extract on DPPH (2,2-diphenyl-1-picryl-hydrazyl-hydrate) radical was assayed using the method of mensor *et al.*<sup>13</sup>. Measurement of superoxide anion scavenging activity was performed based on the method described by Winterbourne *et al.*<sup>14</sup>. Hydroxyl radical scavenging activity was assayed by using Elizabeth and Rao method<sup>15</sup>.

**Table 1: DPPH scavenging potential of ethanolic extract of *Cordia obliqua***

S. No.	Concentration	Percentage of activity	
		Ethanolic extract of <i>Cordia obliqua</i>	Standard (Rutin)
1	125 µg/ml	15.98 ± 0.22	19.44 ± 0.30
2	250 µg/ml	39.67 ± 0.05	36.03 ± 0.07
3	500 µg/ml	54.93 ± 0.02	51.33 ± 0.05
4	1000 µg/ml	65.43 ± 0.06	69.33 ± 0.07

\*All values are expressed as mean ± SEM for three determinations

**Table 2: Superoxide anion radical scavenging activity of ethanolic extract of *Cordia obliqua***

S. No.	Concentration	Percentage of activity	
		Ethanolic extract of <i>Cordia obliqua</i>	Standard (Quercetin)
1	125 µg/ml	34.01 ± 0.40	35.44 ± 0.48
2	250 µg/ml	49.07 ± 0.37	47.88 ± 0.49
3	500 µg/ml	59.97 ± 0.12	61.39 ± 0.18
4	1000 µg/ml	68.54 ± 0.04	62.28 ± 0.09

\*All values are expressed as mean ± SEM for three determinations

**Table 3: Hydroxyl radical scavenging activity of ethanolic extract of *Cordia obliqua***

S. No.	Concentration	Percentage of activity	
		Ethanolic extract of <i>Cordia obliqua</i>	Standard (Ascorbate)
1	125 µg/ml	15.75 ± 0.06	26.87 ± 0.08
2	250 µg/ml	34.03 ± 0.03	30.30 ± 0.05
3	500 µg/ml	64.35 ± 0.01	60.64 ± 0.02
4	1000 µg/ml	65.45 ± 0.01	55.23 ± 0.01

\*All values are expressed as mean ± SEM for three determinations

## RESULTS AND DISCUSSION

### DPPH scavenging activity

DPPH assay are widely used for the evaluation of the antioxidant capacities of natural substances, they both are spectrophotometric approaches based on quenching of stable colored radicals (DPPH) and show the radical scavenging ability of antioxidants even when present in compound biological substances such as plant extracts. The ethanolic extract of *Cordia obliqua* exhibited a maximum DPPH scavenging activity of 65.43% at 1000 µg/ml whereas for Rutin (standard) was found to be 69.33% at 1000 µg/ml. (Table 1)

### Superoxide anion scavenging activity

Superoxide anions destruct bio molecules precisely or indirectly by forming H<sub>2</sub>O<sub>2</sub>, per oxy nitrite or singlet O<sub>2</sub> during aging and pathological measures such as ischemic reoxygenation injury. Superoxide has too been experiential to exactly initiate lipid per oxidation. The maximum scavenging activity of plant extract and Quercetin at 1000 µg/ml was found to be 68.54% at 1000 µg/ml and 62.28% at 1000 µg/ml respectively. Superoxide scavenging ability of plant extract might primarily be due to the presence of flavonoids. (Table 2)

### Hydroxyl radical scavenging activity

Hydroxyl radical is profoundly reactive oxygen centered radical comprise from the response of various hydroperoxides with transition metal ions. It attacks proteins, chromosome, polyunsaturated fatty acid in membranes, and generally biological molecule it contacts and is acknowledged to be gifted of abstracting hydrogen atoms from membrane lipids and brings about per oxidic reaction of lipids. The ethanolic extract of *Cordia obliqua* was exhibited a maximum Hydroxyl radical scavenging activity of 65.45 % at 1000 µg/ml whereas for ascorbate (standard) was found to be 55.23 % at 1000 µg/ml. (Table 3)

## CONCLUSION

In nature there are a wide variety of naturally occurring antioxidants will be available. Antioxidants derived from food and medicinal plants have been increasingly investigated for their various nutritional function and health benefits. Based on the above results, this present study was concluded that the ethanolic extract of aerial parts of *Cordia obliqua* also known as clammy cherry, which contains large amounts of phenolic compounds, exhibits high antioxidant and free radical scavenging activities. These assays indicate that this plant extract is a significant source of natural antioxidant, especially polyphenols and carotenoids, exhibit a wide range of biological effects, including anti-inflammatory, anti-aging, anti-atherosclerosis and anticancer.

## REFERENCES

1. Vivek Kumar Gupta and Surendra Kumar Sharma. Plants as natural antioxidants. Natural Products Radiance 2006; 5 Suppl 4: 326-334.
2. Sunil Kumar. The Importance of antioxidant and their role in pharmaceutical science - A Review. Asian Journal of Research in Chemistry and Pharmaceutical Sciences 2014; 1 Suppl 1: 27 - 44.
3. Parmar C, Kaushal MK. Wild fruits of the Sub-Himalayan region. In: Wild fruits. Kalyani Publishers, New Delhi, India; 1982. p. 19-22.
4. Gupta R, Gupta GD. A review on plant *Cordia obliqua* Willd. (Clammy cherry). Pharmacognosy Reviews 2015; 9 Suppl 18: 127-131.
5. Kritikar KR, Basu BD. Dehradun: Shiva offset press: Indian medicinal plants; 1998. p. 1674-81.
6. Thirupathi K, Kumar SS, Raju VS, Ravikumar B, Krishna DR, Mohan GK. A review of medicinal plants of the genus *Cordia*: Their chemistry and pharmacological uses. Journal of Natural Remedies 2008; 8 Suppl 1: 1-10.
7. Aswal BS, Goel AK, Kulshrestha DK, Mehrotra BN, Patnaik GK. Screening of Indian plants for biological activity: Part XV. Indian Journal of Experimental Biology 1996; 34 Suppl 5: 444-467.

8. Yadav R, Yadav SK. Evaluation of antimicrobial activity of seeds and leaves of *Cordia obliqua Willd* against some oral pathogens. *Indo-American Journal of Pharmaceutical Research* 2013; 3 Suppl 8: 6035-6043.
9. Agnihotri VK, Srivastava SD, Srivastava SK, Pitre S, Rusia K. Constituents from the seeds of *Cordia obliqua* as potential anti-inflammatory agents. *Indian Journal of Pharmaceutical Sciences* 1987; 49 Suppl 2: 66-9.
10. Abou Shaaban RR, Al Angari AA, El Tahir KE, Al Khamis KI, Mirghani OM. Comparative hypotensive and respiratory stimulation effects of ripe and unripe fruit mucilage of *Cordia myxa* and *Cordia obliqua* in guinea pigs and rabbits. *Phytotherapy Research* 1989; 3 Suppl 4: 126-31.
11. Siva Krishnan S, Kottaimuthu A. Hepatoprotective activity of ethanolic extract of aerial parts of *Albizia procera roxb* (*Benth.*) against paracetamol induced liver toxicity on wistar rats. *International Journal of Pharmacy and Pharmaceutical Sciences* 2014; 6 Suppl 1: 233-238.
12. Siva Krishnan S, Kottaimuthu A. Kavitha J. GC-MS analysis of ethanolic extract of aerial parts of *Albizia procera* (*Roxb* *Benth.*) *International Journal of Pharmacy and Pharmaceutical Sciences* 2013; 5 Suppl 3: 702-704.
13. Mensor L, Meneze L, Leitao FS, Reis AS, Dos Santor JC, Coube CS and Leitao SG. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytotherapy Research* 2001; 15 Suppl 2: 127-130.
14. Winterbourne CC, Hawkins RE, Brain M and Carrel RW. The estimation of red cell superoxide dismutase activity. *Journal of Clinical Chemistry and Laboratory Medicine* 1975; 85 Suppl 2: 337-341.
15. Elizabeth K and Rao MNA. Oxygen radical scavenging activity of curcumin. *International Journal of Pharmaceutics* 1990; 58 Suppl 3: 237-240.

**Cite this article as:**

S. Sivakrishnan and M. Swamivelmanickam. *In-vitro* Free Radical Scavenging activity of Aerial parts of ethanolic extract of *Cordia obliqua*. *Int. Res. J. Pharm.* 2019;10(8):102-104  
<http://dx.doi.org/10.7897/2230-8407.1008253>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.