

## PHYTOCHEMICAL STUDIES OF STEM BARK OF *MICHELIA CHAMPACA* LINN

Kodongala Subraya Chandrashekar<sup>1\*</sup>, HegdeVignesh<sup>2</sup> and KodongalaSubrayaPrasanna<sup>3</sup>

<sup>1</sup>Department of Pharmacognosy, Manipal College of Pharmaceutical Sciences, Manipal University, Manipal, Karnataka, India

<sup>2</sup>Department of Pharmacognosy, NGSM Institute of Pharmaceutical Sciences, Paneer, Deralakatte, Mangalore, Karnataka, India

<sup>3</sup>Department of community medicine, Father Muller Medical college, Mangalore, Karnataka, India

\*Chandrashekar Kodongala Subraya, Department of Pharmacognosy, Manipal College of Pharmaceutical Sciences, Manipal University, Manipal-576104 Email: cksbhat@yahoo.co.in

Article Received on: 12/11/10 Revised on: 21/11/10 Approved for publication: 03/12/10

### ABSTRACT

Phytochemical studies of stem bark of *Michelia champaca* Linn ( Family: Magnoliaceae) reveals the presence of Triterpenoids, Steroids, Fatty acid and absence of Alkaloids , Carbohydrates, Flavanoids, Glycosides , Resins, Saponins , Tannins have been reported in this plant for the first time.

**KEY WORDS:** *Michelia champaca*, Magnoliaceae, Triterpenoids, Steroids, Fatty acid

### INTRODUCTION

*Michelia champaca* Linn is a tall and handsome, evergreen tree with a straight trunk; branches ascending, spreading, forming a close head. Belonging to the family Magnoliaceae. It mainly distributed in the sub-Himalayan tract and lower hills up to 3000 ft. in Assam, Burma, Western Ghats, South India. Much cultivated in various parts of India and Burma-Yunnan, Indo-China, Siam, Malaya (not in Malay Peninsula)

The bark is bitter with a sharp acrid taste; causes warmth in the abdomen; the bark is also used as antidote for various poisons; it has vermifuge properties; and facilitates micturition, diuretic, diaphoretic, aphrodisiac; which also removes “kapha” and “vata”, bile and blood affections.

When dried roots and root-bark, mixed with curdled milk, is useful as an application to abscesses, clearing away or maturing the inflammation. In the form of an infusion it is a valuable emmenagogue (Rheede). It is also considered purgative. The bark shows febrifugal properties; it is a stimulant, expectorant, and astringent. Decoction of the bark was tried in sixteen mild cases of chronic gastritis<sup>1</sup>. The present investigation was undertaken to study the Phytochemicals present in the stem bark of the plant.

### MATERIALS AND METHODS

The Stem bark of *Michelia champaca* collected from local area of Mangalore district, Karnataka, India during December 2008. The stem bark was authenticated by Dr. Noeline J. Pinto, Professor and Head of Botany, Dept St. Agnes College Mangalore.

#### Preparation of extracts

The stem bark were sun dried after washing and then grinded to a coarse powder in a grinder. The coarse powder form of stem bark *Michelia champaca* will be exhaustively extracted by soxhlet extractor, using various solvent.

The extract so obtained were subjected to solvent evaporation by vacuum distillation and dried in desiccators. The dried material were tested for different phytoconstituents like Alkaloids, Carbohydrates, Flavanoids, Glycosides, Resins, Saponins , Tannins by standard methods<sup>2-5</sup>.

## REAGENTS

All reagents were of analytical grade and obtained from S.D Fine Chemicals Ltd., Mumbai

## METHODS

### 1. Alkaloids

#### (a) Dragendorff's test

To 2 mg of the ethanolic extract 5 ml of distilled water was added, 2M Hydrochloric acid was added until an acid reaction occurs. To this 1 ml of Dragendorff's reagent was added. Formation of orange or orange red precipitate indicates the presence of alkaloids.

#### (b) Hager's test

To 2 mg of the ethanolic extract taken in a test tube, a few drops of Hager's reagent was added. Formation of yellow ppt confirms the presence of alkaloids.

#### (c) Wagner's test

2 mg of ethanolic extract was acidified with 1.5 % v/v of hydrochloric acid and a few drops of Wagner's reagent was added. A yellow or brown ppt. indicates the presence of alkaloids.

#### (d) Mayer's test

To a few drops of the Mayer's reagent, 2 mg of ethanolic extract was added. Formation of white or pale yellow precipitate. Indicates the presence of alkaloids.

### 2. Carbohydrates

#### (a) Anthrone test

2 mg of ethanolic extract was shaken with 10ml of water, filtered and the filtrate was concentrated. To this 2ml of anthrone reagent solution was added. Formation of green or blue colour indicates the presence of carbohydrates.

#### (b) Benedict's test

2 mg of ethanolic extract was shaken with 10ml of water, filtered and the filtrate was concentrated. To this 5 ml of Benedict's solution was added and boiled for 5 minutes. Formation of brick red coloured ppt indicates the presence of carbohydrates.

#### (c) Fehling's test

2 mg of ethanolic extract was shaken with 10ml of water, filtered and the filtrate was concentrated. To this 1 ml mixture of equal parts of Fehling's solution A and B were added and boiled for few minutes. Formation of red or brick red coloured precipitate indicates the presence of reducing sugar.

#### (d) Molisch's test

2 mg of ethanolic extract was shaken with 10ml of water, filtered and the filtrate was concentrated. To this 2 drops of freshly prepared 20% alcoholic solution of  $\alpha$ - naphthol was added. 2 ml of conc. sulphuric acid was added so as to form a layer below the mixture. Red-violet ring appear, indicating the presence of carbohydrates which disappear on the addition of excess of alkali.

### 3. Flavonoids

#### (a) Shinoda's test

2 mg of ethanolic extract was dissolved in 5ml of ethanol and to this 10 drops of dilute hydrochloric acid followed by a small piece of magnesium were added. Formation of pink, reddish or brown colour indicates the presence of flavonoids.

### 4. Glycosides

#### Molisch's test

2 mg of ethanolic extract was shaken with 10ml of water, filtered and the filtrate was concentrated. To this 2-3 drops of Molisch's reagent was added, mixed and 2ml of concentrated sulfuric acid was added carefully through the side of the test tube. Reddish violet ring appear, indicating the presence of glycosides.

### 5. Triterpenoids

#### Liebermann - Burchard's test

2 mg of dry extract was dissolved in acetic anhydride, heated to boiling, cooled and then 1 ml of concentrated sulphuric acid was added along the sides of the test tube. Formation of a pink colour indicates the presence of triterpenoids.

#### 6. Resins

1 ml of ethanolic extract was dissolved in acetone and the solution was poured in distilled water. Turbidity indicates the presence of resins.

#### 7. Saponins

In a test tube containing about 5 ml of an ethanolic extract, a drop of sodium bicarbonate solution was added. The test tube was shaken vigorously and left for 3 minutes. Formation of honeycomb like froth indicates the presence of saponins.

#### 8. Steroids

##### (a) Liebermann-Burchard's test

2 mg of dry extract was dissolved in acetic anhydride, heated to boiling, cooled and then 1 ml of concentrated sulphuric acid was added along the sides of the test tube. Formation of green colour indicates the presence of steroids.

##### (b) Salkowski reaction

2 mg of dry extract was shaken with chloroform, to the chloroform layer sulphuric acid was added slowly by the sides of test tube. Formation of red colour indicated the presence of steroids.

#### 9. Tannins

To 1-2 ml of the ethanolic extract, few drops of 5% w/v FeCl<sub>3</sub> solution was added. A green colour indicated the presence of gallotannins, while brown colour indicates the presence of pseudotannins.

## RESULTS

The detailed result of the Phytochemical tests carried out on stem bark of the *Michelia chapmpaca* .are presented in Table 1. In this present investigation the phytochemical test reveals the presence of the Triterpenoids and Steroids in petroleum ether extract. Alkaloids ,Carbohydrates, Flavanoids, Glycosides , Resins, Saponins, Tanins are absent in all the five extract.

## REFERENCES

1. Kirtikar KR, Basu BD. Indian Medicinal Plant. New Delhi, Periodical Expert Book Agency; 1991, Vol. 1: 54-59.
2. Johonsen DA, "Plant microtechnique", McGraw-Hill Book Company Inc., New York, 1940, 182
3. Harbone J B, "Phytochemical methods", A guide to modern technique of Plant Analysis, Chapman and Hall, London, 1973
4. Trease, G E and Evans, W C., "Pharmacognosy", 9<sup>th</sup> Edn ELBS Publication, 1985.
5. Anjaneyulu ASR, Jaganmohan KR, Ramachandra LR and Subramanyam C, "Crystalline constituents of euphorbiaceae-XII". Tetrahedron.29,1291-1298.

Table 1: Phytochemical screening of stem bark extract of *Michelia champaca*

<b>Name of extract</b>	<b>Petroleum ether(60-80<sup>o</sup>)</b>	<b>Diethyl ether</b>	<b>Ethyl acetate</b>	<b>Methanol</b>	<b>Water</b>
<b>Alkaloids</b>	-	+ve	-	-	-
<b>Carbohydrates</b>	-	-	-	-	-
<b>Flavanoids</b>	-	-	-	-	-
<b>Glycosides</b>	-	-	+ve	-	-
<b>Triterpenoids</b>	+ve	-	-	-	-
<b>Resins</b>	-	-	-	-	-
<b>Saponins</b>	-	-	-	-	-
<b>Steroids</b>	+ve	-	-		-
<b>Tannins</b>	-	-	-	-	-

Source of support: Nil, Conflict of interest: None Declared