

Research Article



INTERNATIONAL RESEARCH JOURNAL OF PHARMACY

www.irjponline.com

ISSN 2230-8407 [LINKING]

RELATIONSHIP BETWEEN ACNE VULGARIS AND HIRSUTISM TO INSULIN RESISTANCE AND TO ASSESS THE SEVERITY OF TWO CONDITIONS WITH INCREASED INSULIN RESISTANCE

Dr. Mohammed Ibrar Alam

Assistant Professor, Department of Dermatology, Kanti Devi Medical College, Hospital & Research Centre, Mathura, Uttar Pradesh

Address for correspondence

Email: dribraralam@gmail.com

How to cite: Alam MI. Relationship Between Acne Vulgaris And Hirsutism To Insulin Resistance And To Assess The Severity Of Two Conditions With Increased Insulin Resistance. International Research Journal of Pharmacy. 2016;7:2:25-30.

doi:10.7897/2230-8407.07215

=====

ABSTRACT

Background: Acne and hirsutism are common disorders that are simple to cure. They do, however, have a noticeable effect on how one perceives oneself. There is a lot of research on the pathophysiology of acne and hirsutism, and insulin resistance is one of them. They are not fully understood, though.

Aim: The purpose of this study was to determine how insulin resistance is related to acne vulgaris and hirsutism, as well as to evaluate how severe these two disorders are when it comes to insulin resistance.

Methods: A thorough clinical history was taken of each included individual, and then measurements such as height, weight, and BMI were noted, along with any indications or symptoms of underlying insulin resistance. The global acne grading system, or GAGS, was used to grade acne, while the modified Ferriman Gallwey score (mFG) was used to determine the severity of hirsutism. Results were formulated by analysing the data.

Results: Three groups of 174 patients were created, and the group with only acne (154 subjects) showed a strong association ($p < 0.0001$) between acne severity and insulin resistance with recurrence. With twelve participants, the hirsutism-only group had a high association between HOMA-IR and mFG score and recurrence, indicating greater hirsutism severity and increased insulin resistance ($p < 0.0001$ and $p < 0.007$, respectively). Not severity, but BMI and insulin resistance were positively correlated in the combined hirsutism and acne group.

Conclusion: The current study comes to the conclusion that hirsutism and severe, resistant acne are pathophysiologically related conditions, with insulin resistance playing a major role. The psychological well-being of individuals with insulin resistance can be greatly impacted, necessitating the use of an alternative treatment strategy.

Keywords: Acne, global acne grading system, hirsutism, HOMA-IR, mFG

INTRODUCTION

Acne vulgaris and hirsutism are two typical dermatological conditions that are highly prevalent everywhere, especially in India. In Indian circumstances, dermatologists frequently treat these two disorders. Numerous studies have been conducted on the etiopathology and related processes of these entities, which are attributable to a variety of variables. The aetiology of acne vulgaris, a chronic and complicated inflammatory illness originating from the pilosebaceous unit, is multifactorial and includes host factors, genetics, and environmental variables. The etiology is also associated with

increased sebum production, follicular hyperkeratinization, sebaceous gland hyperplasia and hypertrophy, and the growth of *Cutibacterium acnes*.¹

Another dermatological disorder that causes the most anguish in female individuals is hirsutism. Particularly the younger age groups A disorder known as hirsutism is characterised by an abundance of terminal hairs in female patients that resemble the male pattern in locations such as the thighs, upper arms, pelvis, belly, chest, chin, and/or upper lips.²

Numerous ideas have been proposed to explain the pathophysiology of acne vulgaris and hirsutism. The function of insulin resistance (IR), which has lately been acknowledged and investigated by several physicians and academics, is one such notion. A hyperglycemic condition brought on by insulin resistance and a reduction in the insulin-responsive tissues' ability to absorb glucose from circulation at the same insulin concentration has detrimental effects on both macro and microvascular health. The prevalence of insulin resistance has significantly grown due to sedentary lifestyles and contemporary technology.³

Insulin resistance is linked to a number of dermatological disorders, such as vitiligo vulgaris, androgenetic alopecia, psoriasis, and acanthosis nigricans, in addition to hirsutism and acne. Additionally, there is a correlation between insulin resistance and several metabolic illnesses such as dyslipidemia, diabetes mellitus, and PCOS (polycystic ovarian syndrome) as well as ischemic heart disease. In addition to these negative effects, hirsutism and acne vulgaris in afflicted individuals have a major psychological influence.⁴

The current clinical investigation sought to determine the correlation between insulin resistance and hirsutism and acne vulgaris, as well as the degree of these two disorders. The current cross-sectional observational study was out to evaluate the severity of two conditions with elevated insulin resistance and to determine a link between hirsutism and acne vulgaris and insulin resistance.

MATERIALS AND METHODS

The study was carried out at the dermatology, venereology, and leprosy departments following approval from the relevant institutional ethical committee. The research population was drawn from the institute's Department of Dermatology. Study participants must meet the inclusion and exclusion criteria for research participation in addition to being hirsutism and acne vulgaris patients who visit the outpatient department of dermatology. Prior to participation, informed permission was obtained in writing and verbally from every study participant. Subjects with systemic inflammatory diseases affecting insulin metabolism, hormonal therapy, history of oral retinoid use, use of alternative medications, pregnancy, lactation, malignancy, atherosclerotic vascular diseases, hypertension, and thyroid dysfunction were excluded from the study.

Each subject had a comprehensive history taking after final inclusion, which was succeeded by a clinical examination. All of the individuals' demographic information, including their height, weight, and body mass index, was entered onto a pre-made, organised proforma that also evaluated any symptoms or indicators of underlying insulin resistance. Recurrent infections of the bladder, vagina, skin, or gums; irregular menstruation; deepening of the voice; excessive hair loss; seborrhea; decreased or increased appetite; weakness or exhaustion; and quick or unexplained weight loss or gain were among the concerns on the proforma.

The global acne grading system, or GAGS, was used to grade the severity of acne, while the modified Ferriman Gallwey (mFG) scores were used to evaluate the severity of hirsutism.

In order to evaluate the statistical relationship between hirsutism, acne, and insulin resistance, the collected data were statistically analysed using SPSS software version 21.0 and the Pearson correlation coefficient. The data's confidentiality was maintained at all times. When the p-value was less than 0.05, it was deemed statistically significant.

RESULTS

Of the 174 people in the research, 3 groups were identified: Group I had 88.50% (n=154) of subjects with acne, Group II had 6.89% (n=12) of respondents with hirsutism, and Group III had 4.59% (n=8) of subjects with both hirsutism and acne vulgaris, respectively. The research participants were 24.81 ± 4.32 years old on average.

The majority of research participants had normal BMIs, with 76 having mean BMIs of 24.26 and BMIs ranging from 18.5 to 24.9. A noteworthy correlation with insulin resistance was seen in 41.4% (n=32) of the participants. Eleven research participants from group I with a BMI of greater than thirty showed a 100% high correlation with insulin

resistance. Out of 40 participants, 46 individuals with a BMI between 25 and 30 who were overweight also exhibited insulin resistance. Out of the 21 participants, BMI <18.5 underweight category, 2 were exhibited insulin resistance. The study discovered that there were significant elevations in the mean fasting blood sugar, mean fasting insulin, and mean HOMA-IR. Age, recurrence, increased HOMA-IR, and GAGS all showed a substantial ($p<0.01$) correlation with BMI in this cohort.

Age and BMI were shown to be significantly correlated ($p<0.05$). However, when employing Pearson's correlation coefficient, as shown in Table 1, no significant link was found between BMI and gender with $p>0.05$. The groups with the highest percentages of insulin resistance were those with very severe acne (59%), severe acne (88%), and moderate acne (46%), respectively.

Seven of the twelve female individuals in group II who had a verified diagnosis of hirsutism were found to have insulin resistance.

Of the seven participants with insulin resistance, five were naïve, whereas the other three had recurrent hirsutism. The mean BMI of the hirsute patients was 24.26. One patient with a BMI of more than thirty had insulin resistance. Out of the five participants, four had a high BMI in the overweight range of 25–30, and three had a normal BMI. These subjects did not exhibit insulin resistance.

Using Pearson's correlation coefficient, age was the only significant factor connected with high BMI ($p<0.05$). Of the study individuals with high BMI, 25% ($n=3$) were between the ages of 21 and 30. In contrast, 20.8% ($n=3$) of the study subjects with high BMI were between the ages of 31 and 40.

In this group, recurrence, age, and a high HOMA-IR score of >2.5 were all linked with a high BMI ($p<0.01$). Table 2 indicates that no significant connection was seen in mFG scores with $p>0.05$. Eight of the female participants in group III exhibited both acne vulgaris and hirsutism. Of the eight participants, six had just received a diagnosis of hirsutism and acne, whereas two had recurring illness. One of the two patients with recurrent illness was insulin resistant. Three of the seven patients were insulin resistant. Subjects with a mean BMI of 26.6 who were insulin resistant had noticeably higher BMIs. Pearson's correlation revealed no link between elevated mFG or higher GAGS scores and BMI in eight patients in group III.

DISCUSSION

The current cross-sectional observational study was out to evaluate the severity of two conditions with elevated insulin resistance and to determine a link between hirsutism and acne vulgaris and insulin resistance. Of the participants in Group I, II, and III, 14.94% ($n = 26$) had a positive history of type 2 diabetes mellitus, compared to 14.28% ($n = 22$), 16.6% ($n = 2$), and 25% ($n = 2$) study subjects, respectively. Twenty female and 64 male participants showed positive indications and symptoms of insulin resistance.

Insulin resistance was observed in 27 research participants out of 78 patients with naïve acne, and in 76 subjects with recurrent acne, insulin resistance was observed in 57 study participants, suggesting a substantial potential of association. These findings aligned with research conducted in 2021 by Gayen R et al. and in 2012 by Ray S et al., both of whom found comparable demographics among their study participants.

The bulk of research participants, or 76 individuals, had a mean BMI of 24.26 and a BMI in the range of 18.5 to 24.9, according to the study data. A noteworthy correlation with insulin resistance was seen in 41.4% ($n=32$) of the participants. Eleven research participants from group I with a BMI of greater than thirty showed a 100% high correlation with insulin resistance. Out of 40 participants, 46 individuals with a BMI of 25–30 and overweight were insulin resistant. Out of the twenty-one participants in the BMI <18.5 underweight category, two exhibited insulin resistance. These findings corroborated those of Gonzalaz-Gonzalaz J et al⁸ in 2016 and Balato N et al⁷ in 2014, who observed a comparable relationship between BMI and insulin resistance. The study revealed that there was a substantial increase in mean HOMA-IR, mean fasting insulin, and mean fasting blood sugar. In this group, recurrence, age, GAGS, and elevated HOMA-IR all strongly correlated with BMI ($p<0.01$).

Age and BMI were shown to be significantly correlated ($p<0.05$). However, when utilising Pearson's correlation coefficient to analyse the relationship between BMI and gender, no significant link was found with $p>0.05$. The groups with the highest percentages of insulin resistance were those with very severe acne (59%), severe acne (88%), and

moderate acne (46%), respectively. These results were consistent with research conducted by Bloomgarden Z et al. (2009) and Lynn D et al. (2016), who found a similar relationship between BMI and a number of other characteristics.

The results of the study showed that insulin resistance was present in seven out of the twelve female subjects in group II who had a confirmed diagnosis of hirsutism. Of the seven participants with insulin resistance, five were naïve, whereas the other three had recurrent hirsutism. The mean BMI of the hirsute patients was 24.26. One patient with a BMI of more than thirty had insulin resistance. Out of the five participants, four had a high BMI in the overweight range of 25–30, and three had a normal BMI. These subjects did not exhibit insulin resistance. These findings were consistent with earlier research by Abdelmawla M et al. (2011) in 2019 and Collier C et al. (2008), which found a comparable relationship between hirsutism and BMI.

By evaluating the data using Pearson's correlation coefficient, it was also shown that age was the sole significant factor connected with high BMI ($p < 0.05$). Of the study individuals with high BMI, 25% ($n=3$) were between the ages of 21 and 30. In contrast, 20.8% ($n=3$) of the study subjects with high BMI were between the ages of 31 and 40. In this group, there was a significant p -value correlation between a high BMI, a high HOMA-IR score of >2.5 , recurrence, and age. Nevertheless, there was no discernible correlation between the mFG scores and $p > 0.05$. These findings were in line with earlier research by Munichandruppa P et al. (2017) and Bhate K et al. (2013), who revealed a substantial correlation between BMI and HOMA-IR and age.

It was seen that in group III having both hirsutism and acne vulgaris, 8 female subjects had hirsutism and acne. Among these 8 subjects, 6 subjects had a new and recent diagnosis of hirsutism and acne, whereas, recurrent disease was seen in 2 subjects. One of the two patients with recurrent illness was insulin resistant. Three of the seven naïve patients were insulin resistant. With a mean BMI of 26.6, participants with insulin resistance had a considerably higher BMI. Pearson's correlation revealed no link between elevated mFG or higher GAGS scores and BMI in eight patients in group III. Furthermore, no association ($p > 0.05$) was seen between higher severity of acne or hirsutism and insulin resistance.

CONCLUSION

Taking into account its limitations, the current study comes to the conclusion that insulin resistance plays a significant role in the pathophysiology of hirsutism and severe, resistant acne. The psychological well-being of individuals with insulin resistance can be greatly impacted, necessitating the use of an alternative treatment strategy.

REFERENCES

1. Kim J. Review of the innate immune response in acne vulgaris: Activation of toll-like receptor 2 in acne triggers inflammatory cytokine responses. *Dermatology* 2005;211:193-8.
2. Casey J. Hirsutism: Pathogenesis and treatment. *Aust N Z J Med* 1980;10:240-5.
3. Gayen R, Podder I, Chakraborty I, Chowdhury S. Sex hormones, metabolic status, and obesity in female patients with acne vulgaris along with clinical correlation: An observational cross-sectional study. *Indian J Dermatol* 2021;66:60-6.
4. Messenger AG, Sinclair RD, Farrant P, de Berker DAR. Hirsutism. In: Rook A, Burns T, editors. *Rook's Textbook of Dermatology*. Chichester, West Sussex, UK: Wiley-Blackwell; 2010. p. 89.64
5. Ünlühizarci K, Karababa Y, Bayram F, Kelestimur F. The Investigation of Insulin Resistance in Patients with Idiopathic Hirsutism. *J Clin Endocrinol Metab* 2004;89:2741-4.
6. Ray S, Bairagi A, Guha S, Ganguly S, Ray D, Basu A, *et al.* A simple way to identify insulin resistance in nondiabetic acute coronary syndrome patients with impaired fasting glucose. *Indian J Endocrinol Metab* 2012;16:460-4.
7. Balato N, Megna M, Palmisano F, Patrino C, Napolitano M, Scalvenzi M, *et al.* Psoriasis and sport: A new ally? *J Eur Acad Dermatol Venereol* 2014;29:515-20.
8. González-Saldivar G, Rodríguez-Gutiérrez R, Ocampo-Candiani J, González-González J, Gómez-Flores M. Skin manifestations of insulin resistance: From a biochemical stance to a clinical diagnosis and management. *Dermatol Ther* 2016;7:37-51

9. Bloomgarden Z. American Association of Clinical Endocrinologists (AACE) Consensus Conference on the Insulin Resistance Syndrome. *Diabetes Care* 2003;26:1297-303.
10. Lynn D, Umari T, Dellavalle R, Dunnick C. The epidemiology of acne vulgaris in late adolescence. *Adolesc Health Med Ther* 2016;7:13-25.
11. Abdelmawla M, Esawy A, Khater E, Khalifa N. Insulin resistance in androgenetic alopecia and acne vulgaris. *Egypt J Dermatol Venerol* 2019;39:83.
12. Collier C, Harper J, Cantrell W, Wang W, Foster K, Elewski B, *et al.* The prevalence of acne in adults 20 years and older. *J Am Acad Dermatol* 2008;58:56-9.
13. Munichandrappa P, K G M, C K, Variyar A. A comparative study of insulin resistance in acne vulgaris. *Int J Res Dermatol* 2017;3:403.
14. Bhate K, Williams H. Epidemiology of acne vulgaris. *Br J Dermatol* 2013;168:474-85.
15. Wankhade V, Shah V, Tomar S, Singh R. Clinical and investigative study of Hirsutism. *J Clin Diagn Res* 2019;13:1-6.
16. Nagpal M, De D, Handa S, Pal A, Sachdeva N. Insulin resistance and metabolic syndrome in young men with acne. *JAMA Dermatol* 2016;152:399-404.

TABLES

	Recurrent/Naive	Gender	Age	GAGS	HOMA-IR	BMI
Recurrent/Naive						
Pearson correlation	1	-0.016	0.002	0.363	0.277	0.384
P	-	0.73	0.000	0.000	0.000	
N	154	154	154	154	154	154
Gender						
Pearson correlation	-0.016	1	-0.037	0.001	0.055	-0.076
P	0.73	-	0.47	0.93	0.34	0.15
N	154	154	154	154	154	154
Age						
Pearson correlation	0.004	-0.037	1	0.093	0.101	0.184
P	0.95	0.47	-	0.08	0.05	0.001
N	154	154	154	154	154	154
GAGS						
Pearson correlation	0.34	0.001	0.07	1	0.749	0.625
P	0.000	0.73	0.07	-	0.000	0.000
N	154	154	154	154	154	154
HOMA-IR						
Pearson correlation	0.277	0.037	0.101	0.749	1	0.812
P	0.000	0.319	0.068	0.000	-	0.000
N	154	154	154	154	154	154
BMI						
Pearson correlation	0.384	0.076	0.184	0.625	0.812	1
P	0.000	0.171	0.001	0.000	0.000	-
N	154	154	154	154	154	154

Table 1: Pearson’s correlation of various parameters in study subjects with acne vulgaris

	Recurrent/Naive	mFG score	GAGS	HOMA-IR	BMI	Age
Recurrent/Naive						
Pearson correlation	1	0.074	-0.064	0.176	0.317	0.269
p-value	-	0.786	0.815	0.505	0.217	0.302
Number (n)	8	8	8	8	8	8
mFG score						

Pearson correlation	0.074	1	-0.137	-0.248	0.237	-0.197
p-value	0.786	-	0.597	0.343	0.367	0.455
Number (n)	8	8	8	8	8	8
GAGS						
Pearson correlation	-0.064	-0.137	1	0.463	0.259	0.487
p-value	0.815	0.597	-	0.061	0.316	0.047
Number (n)	8	8	8	8	8	8
HOMA-IR						
Pearson correlation	0.176	-0.248	0.463	1	0.519	0.586
p-value	0.501	0.343	0.061	-	0.032	0.012
Number (n)	8	8	8	8	8	8
BMI						
Pearson correlation	0.317	0.237	0.259	0.519	1	0.157
p-value	0.217	0.367	0.316	0.032	-	0.551
Number (n)	8	8	8	8	8	8
Age						
Pearson correlation	0.269	-0.197	0.487	0.586	0.157	1
p-value	0.302	0.455	0.047	0.012	0.555	-
Number (n)	8	8	8	8	8	8

Table 3: Pearson's correlation of various parameters in study subjects with hirsutism and acne vulgaris both